

Assessment of PM_{10} Atmospheric Particle Concentrations and Heavy Metals Levels in the Air of Gorgan City, Iran

Hadi Rahimzadeh ^{1,2} , Hamidreza Heidari ^{1,3} , Somayeh Beirami ^{1,2} , Abdollah Dargahi ^{4,5*} , Hossein Faraji ⁶ , Farnam Mokhtarkoureh ⁷ , Mohammad Heidari ⁸ , Fatemeh Piran ² , Farshad Bahrami Asl ⁹ 

¹ Environmental Health Research Center, Golestan University of Medical Sciences, Gorgan, Iran.

² Department of Environmental Health Engineering, Faculty of Health, Golestan University of Medical Sciences, Gorgan, Iran.

³ Department of Occupational health, Faculty of Health, Golestan University of Medical Sciences, Gorgan, Iran.

⁴ Department of Environmental Health, Khalkhal University of Medical Sciences, Khalkhal, Iran.

⁵ Social Determinants of Health Research Center, School of Health, Ardabil University of Medical Sciences, Ardabil, Iran.

⁶ Tropical and Communicable Diseases Research Center, Iranshahr University of Medical Sciences, Iranshahr, Iran.

⁷ Student of the Research Committee, Khalkhal University of Medical Sciences, Khalkhal, Iran.

⁸ Department of Epidemiology, School of Medicine, Urmia University of Medical Sciences, Urmia, Iran.

⁹ Department of Environmental Health Engineering, School of Public Health, Urmia University of Medical Sciences, Urmia, Iran.

ARTICLE INFO

ORIGINAL ARTICLE

Article History:

Received: 17 May 2026

Revised: 13 August 2026

Accepted: 20 August 2026

*Corresponding Author:

Abdollah Dargahi

Email:

a.dargahi29@yahoo.com

Tel:

+98 914 159 7607

Keywords:

Heavy Metals,

PM_{10} ,

Ambient Air Quality,

Seasonal Variation.

ABSTRACT

Introduction: With increasing industrialization and urbanization, particulate matter (PM_{10}) has become one of the major air pollutants and a critical concern for environmental and public health. Therefore, this study aimed to assess seasonal/annual variations of PM_{10} -bound heavy metals in Gorgan, compare them with WHO/international data, and identify potential sources via correlation analysis.

Methods: cross-sectional experimental study was conducted between March 2023 and March 2024 in Gorgan, Golestan province, Iran. Sampling was performed every six days throughout the study period according to the U.S. Environmental Protection Agency (EPA) method. A total of 52 samples were collected to determine PM_{10} and heavy metals concentrations. Samples were taken from a central urban station using a BGI Omni FT ambient air sampler.

Results: Results showed that PM_{10} concentrations ranged from a minimum of $20.83 \mu g/m^3$ in early November to a maximum of $277.78 \mu g/m^3$ in early February, with a mean $\pm$ SD of $69.3 \pm 49.82 \mu g/m^3$ across the study period. Seven trace elements were analyzed, and their annual mean concentrations followed the order: $Cd < Co < Cr < Ni < Cu < Pb < Zn$. Strong positive correlations were observed among several metals, particularly Cu, Cr, and Ni, with nearly linear relationships.

Conclusion: The findings demonstrated that PM_{10} levels in Gorgan slightly exceeded the WHO 24-hour guideline, with more pronounced fluctuations on a monthly scale. Among the heavy metals, Zn, Pb, and Cu exhibited the highest concentrations, peaking during winter.

Citation: Rahimzadeh H, Heidari H, Beirami S, et al. *Assessment of PM_{10} Atmospheric Particle Concentrations and Heavy Metals Levels in the Air of Gorgan City, Iran*. J Environ Health Sustain Dev. 2026; 11(3): 3127-40.

Introduction

Climate change has been identified by the World Health Organization (WHO) as one of the greatest

health threats of the twenty-first century. Alongside this global issue, air pollution remains a major environmental determinant of public health

and economic stability in many societies¹. The severity of air pollution is largely associated with the absence of bypass infrastructure, proximity of industrial facilities to residential neighborhoods, and extensive use of fuels with low combustion quality^{2, 3}. In addition, poor air circulation intensifies the accumulation of hazardous pollutants such as particulate matter (PM), volatile organic compounds (VOCs), nitrogen oxides (NO_x), carbon monoxide (CO), heavy metals, and persistent organic pollutants¹.

Particulate matter (PM) is a critical air pollutant that adversely affects environmental integrity and human health⁴. PM arises from a range of natural and human-induced sources. Volcanic eruptions, wildfires, dust events, and sea-salt aerosols contribute naturally, while industrial operations, combustion processes, vehicle exhaust, and tobacco smoke are key anthropogenic contributors^{5, 6}. Based on their size, PM is commonly classified into three major categories: coarse, fine, and ultrafine particles. Exposure to suspended particulate matter (SPM) has been linked to a wide range of short- and long-term health effects, including cardiovascular diseases, pulmonary disorders, eye irritation, headaches, and cancers, ultimately contributing to premature mortality⁷. Determining the atmospheric concentration of particles with an aerodynamic diameter $\leq 10 \mu\text{m}$ (PM₁₀) is crucial not only because of their detrimental health effects but also for developing strategies to reduce air pollution. The toxicity of suspended particles depends on both their size and chemical composition; however, the major health risks are primarily attributed to their chemical constituents^{8, 9}. Particulate matter (PM) may contain heavy metals such as lead (Pb), cadmium (Cd), nickel (Ni), copper (Cu), chromium (Cr), manganese (Mn), vanadium (V), zinc (Zn), and arsenic (As), all of which pose serious risks to human health. The principal exposure pathways for these contaminants are inhalation, ingestion, and skin contact⁴. Major contributors to atmospheric metal contamination include smelting processes, fossil fuel burning, mining activities, and road traffic emissions, with particularly high impacts

observed in urban and industrialized areas^{10, 11}. The environmental persistence of heavy metals constitutes a major concern, as they remain stable for extended periods and bioaccumulate in multiple environmental compartments—air, soil, water, and living tissues. Chronic exposure, even at trace levels, may lead to disturbances in biological functioning¹².

Given the detrimental effects of heavy metals on both human health and ecosystems, numerous investigations have been carried out globally to quantify their environmental levels^{9, 13}. Arsenic, cadmium, and nickel have been classified by the International Agency for Research on Cancer (IARC) as Group 1 human carcinogens, with established associations with lethal cancers of the lung, trachea, and bronchi¹. Lead has been categorized as a probable human carcinogen (Group 2A) and is known to cause neurological and renal damage¹⁴. Chronic exposure in children has been associated with developmental delays and deficits in learning¹⁵. Elevated airborne copper (Cu), frequently used in industry, is associated with oxidative stress and liver damage¹⁶. In general, exposure to heavy metals may disrupt neurological function, impair renal health, interfere with bone development, and cause damage to various other organs^{17, 18}. Recently, the European Parliament and Council proposed updated guidelines for ambient air quality and cleaner air in Europe. According to this proposal, the annual limits for PM₁₀ are reduced from 40 $\mu\text{g}/\text{m}^3$ to 20 $\mu\text{g}/\text{m}^3$, and for PM_{2.5} from 25 $\mu\text{g}/\text{m}^3$ to 10 $\mu\text{g}/\text{m}^3$, with the introduction of a new daily limit for PM_{2.5}¹⁹. Risk-based evaluation of toxic metal exposure in indoor and outdoor environments across China revealed that zinc recorded the highest concentrations, whereas lead showed the lowest. Notably, the lifetime cancer risk (LCR) estimates for arsenic and cadmium were above the acceptable limits in specific cities²⁰. Additionally, a study examining the impact of PM₁₀ on mortality in Bogotá from 1998 to 2006 found that increases in PM₁₀ concentrations were associated with acute respiratory-related deaths²¹. More recently, two studies reported that PM₁₀ levels in Santa Marta

exceeded the international precautionary threshold of 50 µg/m³ ²¹. Despite the growing body of research on urban air pollution in Iran, comprehensive data on PM₁₀ and its associated heavy metal content remain scarce for many mid-sized cities, particularly those with distinct climatic and geographical characteristics. Gorgan, the capital of Golestan Province, exemplifies such a city. It features a temperate climate with high humidity in summer and cold winters—conditions that can substantially affect the atmospheric dispersion, transformation, and deposition of particulate-bound pollutants. Moreover, rapid urbanization and increased vehicular traffic in recent years have heightened concerns about local air quality, yet no systematic monitoring or characterization of PM₁₀-bound trace elements has been conducted in this region. This study addresses this critical knowledge gap by providing the first year-round dataset on PM₁₀ and its associated heavy metals (Zn, Cu, Cr, Cd, Pb, Ni, and Co) in Gorgan's urban atmosphere. Accordingly, the specific objectives of this investigation were to: (1) determine the seasonal and annual concentration profiles of PM₁₀ and the selected heavy metals; (2) compare the measured concentrations with World Health Organization (WHO) air quality guidelines and findings from comparable international studies; and (3) evaluate inter-element correlations to identify potential common sources. We hypothesized that heavy metal concentrations would exhibit significant seasonal variations, with peak levels in winter driven by increased fossil fuel combustion, and that strong positive correlations among specific metals would indicate shared anthropogenic emission sources, particularly traffic-related and industrial activities.

Materials and Methods

Study Area

This experimental cross-sectional study was conducted between March 2023 and March 2024 in Gorgan, the capital of Golestan Province, located in northeastern Iran. Gorgan is situated at latitude 36.50°N and longitude 54.26°E, at an altitude of 160 meters above sea level. The city has

a temperate climate characterized by hot, humid summers and cold winters. Gorgan comprises three distinct topographical regions: plains, piedmonts, and mountains. The prevailing wind direction is from the west and southwest. According to the most recent census, the population of Gorgan is approximately 500,000 inhabitants.

Sampling and Analytical Procedures

Air sampling was conducted on a weekly basis (once every 7 days) over a 12-month period from March 2023 to March 2024, yielding a total of 52 samples. Samples were collected at a fixed monitoring station located in the city center, using a BGI OMNI FT ambient air sampler operating at a constant flow rate of 5 L/min, in accordance with the U.S. Environmental Protection Agency (EPA) reference method for PM₁₀ ²².

Particulate matter (PM₁₀) was collected on quartz microfilters. Before and after sampling, all filters were conditioned in a desiccator for at least 24 h to eliminate moisture. The filters were weighed using a microbalance with 0.001 mg precision, and PM₁₀ mass concentrations (µg/m³) were calculated gravimetrically based on the difference in filter weights ($W_2 - W_1$).

For heavy metal analysis, the loaded filters were digested using a mixture of nitric acid and perchloric acid in a 10:1 (v/v) ratio, following NIOSH Method 7300 (23). The digested solutions were filtered and stored in polyethylene bottles under refrigeration (4°C) until analysis. Heavy metal concentrations (Zn, Cu, Cr, Cd, Pb, Ni, and Co) were determined using voltammetry with a 797 VA Computrace polarograph (Metrohm, Switzerland).

Quality control measures included the following: (i) the use of field blanks (one blank per 10 samples) to correct for potential contamination during handling and transport; (ii) pre-baking of quartz filters at 450°C for 4 hours to remove organic impurities; and (iii) calibration of the sampler before each sampling campaign using a certified flow meter. The limit of detection (LOD) and limit of quantification (LOQ) for each heavy metals were calculated as 3 and 10 times the

standard deviation of blank measurements, respectively. The LOD values (in ng/m³) were: Zn: 0.5, Cu: 0.2, Cr: 0.1, Cd: 0.05, Pb: 0.3, Ni: 0.15, and Co: 0.08. Concentrations below the LOD were replaced with LOD/√2 for statistical analysis.

Statistical Analysis

Quantitative data are expressed as mean ± standard deviation (mean ± SD). Prior to parametric testing, the normality of data distribution was assessed using the Shapiro–Wilk test. All variables met the normality assumption ($p > 0.05$), justifying the use of parametric tests. One-way analysis of variance (ANOVA) was used to compare PM₁₀ and heavy metal concentrations across seasons. When ANOVA indicated significant differences ($p < 0.05$), Tukey's HSD post-hoc test was applied for pairwise comparisons. Pearson correlation coefficients were calculated to evaluate linear relationships between variables. All statistical analyses were performed using STATA version 18 (StataCorp, USA). Graphical visualizations, including correlation matrices and scatterplots, were generated using R

software (version 4.2) with the 'ggplot2' package. Statistical significance was set at a two-tailed p -value < 0.05 for all analyses.

Results

PM₁₀ Concentrations

To assess pollution levels and their association with nearby dust sources, the variations in PM₁₀ and heavy metal concentrations in Gorgan were analyzed over a 12-month period. The results showed that PM₁₀ concentrations ranged from a minimum of 20.83 µg/m³ in early November to a maximum of 277.78 µg/m³ in early February, with a mean ± standard deviation of 69.3 ± 49.82 µg/m³ over the study period (Table 1). These values are slightly higher than the World Health Organization's 24-hour guideline of 50 µg/m³. Seasonal mean concentrations ranged from 65.97 ± 36.72 µg/m³ in spring to 71.58 ± 50.34 µg/m³ in summer, with no statistically significant differences observed between the seasons ($p = 0.992$) (Table 2). Monthly variations showed the highest concentration in July (114.58 µg/m³) and the lowest in January (43.40 µg/m³) (Figure 1).




Figure 1: Monthly variations of PM₁₀ concentrations in Gorgan.

Table 1: Summary comparison of PM₁₀ concentrations obtained in the present study with those reported in other studies conducted in Iran and other countries (expressed in µg/m³).

Ref	Duration	PM ₁₀	Place
Current study	March 2023 to March 2024	69.3	Gorgan
Aldeen et al. 2025 ²³	October and November 2023	85.11 72.82	the National Iraqi Museum The Natural History Museum Baghdad
Adly et al. 2025 ²⁴	December 2020 to December 2024	127.7 130.2 132.9 135.5 138.3	Saudi Arabia (Makkah)
Kebe et al. 2025 ⁴	In 2018 and 2019	232.318 209.854	Hlm (industrial site) Yoff (coastal and urban site) Senegal
Shahsavani et al. 2025 ²⁰	summer of 2019	78.12	iran(Shiraz)
	2008–2019	223.4	
	January	177.09	
	February	265.19	
	March	179.08	
	April	192.43	
	May	192.43	
Hamidi et al. 2025 ²⁵	June	346.9	iran(Ahvaz)
	July	388.18	
	August	189.59	
	September	188.04	
	October	167.46	
	November	143.77	
	December	148.97	
Alharbi et al. 2024 ²⁶	April–December2023	223.12	Saudi Arabia(Riyadh)
Zeb et al. 2024 ²⁷	December 2017 to February 2018 (winter)	586	Pakistan(Khyber Pakhtunkhwa)
Lara et al. 2023 ²⁸	May 2019 to September 2020	31	Spain(Gijon)
Ahmad et al. 2023 ²⁹	winter 2019	279.73	Pakistan (Lahore)
Esform et al. 2022 ³⁰	September 2019 to March 2020	97.5	Iran(Birjand)
Zeb et al. 2022 ³¹	winterseason(2018)	505.06 255.04 238.48	industrial urban suburban Pakistan(Nowshera)
Sirithian et al. 2022 ³²	(January to April) from 2016 to 2020.	73.86 64.58	urban rural Thailand
Usman et al. 2022 ³³	year of 2019	64	Pakistan(Hindu KushRegion)
Kliengchuay et al. 2021 ³⁴	July 2009 to December 2017	42.98	Thailand (Lamphun)
Liu et al. 2020 ³⁵	February to March 2016	98	China(Tian'jin)
Gupta et al. 2020 ³⁶	January 2013 to December 2018	136.25	Bangladesh
Shahid et al.	March and April of 2010	438	Pakistan (Karachi)

Ref	Duration	PM ₁₀	Place
2018 ³⁷			
Farsani et al. 2018 ³⁸	normal days	autumn	131
		winter	189
		spring	145
		summer	127 µg
	Duststorm	autumn	410
		winter	742
		spring	300
		summer	278
Di Vaio et al. 2018 ³⁹	February 2013- October 2013		
	autumn	30.22	Italy (Acerra)
	winter	39.6	
	spring	33.85	
	summer	35.46	
Munir et al. 2017 ⁴⁰	January 2014 to September 2015	44.53	Saudi Arabia (Makkah)
Alam et al. 2014 ⁴¹	March and April of 2010	406.2	Pakistan (Lahore)
Sharma et al. 2014 ⁴²	January 2010 to December 2011	191.4	India (Delhi)

Table 2: Mean and standard deviation of PM₁₀ and heavy metal concentrations by season in Gorgan City.

	Season					Test
	Spring	Summer	Autumn	Winter	Total	
PM ₁₀	65.972 (36.721)	71.581 (50.348)	71.047 (41.778)	68.866 (73.818)	69.311 (50.312)	0.992
Zn	226.714 (75.242)	277.385 (108.586)	461.538 (223.071)	541.500 (96.514)	370.731 (186.583)	< 0.001
Cu	13.532 (5.625)	7.507 (2.668)	27.999 (8.825)	40.082 (5.227)	21.769 (13.901)	< 0.001
Cd	2.529 (0.823)	2.438 (0.747)	3.500 (0.998)	4.300 (1.437)	3.158 (1.250)	< 0.001
Pb	82.286 (33.160)	56.000 (26.331)	157.846 (54.714)	178.667 (39.204)	116.846 (63.864)	< 0.001
Cr	6.720 (2.374)	4.046 (1.911)	9.359 (2.675)	13.009 (1.755)	8.162 (3.928)	< 0.001
Co	3.953 (1.397)	2.333 (1.147)	4.417 (1.099)	5.955 (0.844)	4.126 (1.697)	< 0.001
Ni	13.439 (4.748)	8.091 (3.822)	18.718 (5.349)	26.019 (3.511)	16.325 (7.856)	< 0.001

Heavy Metal Concentrations

Analysis of seven trace elements in ambient air samples yielded their annual mean concentrations, as shown in Table 3, with comparative data from other studies presented in Table 4. The mean concentrations followed the order: Cd (3.2 ± 1.23 ng/m³) < Co (4.1 ± 1.68) < Cr (8.2 ± 3.88) < Ni (16.3 ± 7.77) < Cu (21.8 ± 13.76) < Pb (116.8 ± 63.24) < Zn (370.7 ± 184.78). All metals exhibited considerable seasonal variations, with statistically

significant differences ($p < 0.001$). Zinc (Zn) showed a continuous increasing trend from spring (226.7 ng/m³) to winter (541.5 ng/m³). For the other metals, the minimum concentrations were observed in summer, while the maximum concentrations were observed in winter. The ratio of seasonal variations (minimum/maximum) ranged from 1.76 for Cd to 5.34 for Cu, indicating a marked increase in metal concentrations during winter (Table 2).

Table 3: Annual mean concentrations of heavy metals in PM₁₀ (ng/m³).

	Zn	Cu	Cd	Pb	Cr	Co	Ni
mean	370.7 ± 184.78	21.8 ± 13.76	3.2 ± 1.23	116.8 ± 63.24	8.2 ± 3.88	4.1 ± 1.68	16.3 ± 7.77
min	150.0	2.8	1.2	24.0	1.9	1.1	3.7
max	870.0	51.7	6.1	240.0	16.7	7.4	33.3

Table 4: Summary comparison of heavy metal concentrations associated with PM₁₀ obtained in the present study and those reported in other studies conducted in Iran and other countries (ng/m³)..

Ref	Duration	Zn	Cu	Cd	Pb	Cr	Co	Ni	Location
Shahsavani et al. 2025 ²⁰	summer of 2019	33.08	102.04	–	17.61	–	–	82.15	Iran (Shiraz)
Alharbi et al. 2024 ²⁶	April–December 2023	9.06	10.97	0.04	1.11	3.75	0.25	1.72	Saudi Arabia (Riyadh)
Lara et al. 2023 ²⁸	May 2019 to September 2020	9.4	9.2	0.19	9.4	3.2	0.21	1.7	Spain (Gijon)
Karimi et al. 2023 ⁴³	2021 autumn	29.21	–	–	6/81	29/55	–	–	Iran (Tehran)
Esform et al. 2022 ³⁰	September 2019 to March 2020	0.71	0.006	9.588	0.004	0.006	0.224	0.006	Iran (Birjand)
Rezayani et al. 2022 ⁴⁴	April 2019 to March 2020	106.21	16.377	5.357	31.65	5.615	–	5.77	Iran (Tehran)
Karimi et al. 2022 ⁴⁵	2021 spring	18.78	–	–	0/17	25/42	–	–	Iran (Tehran)
Karimi et al. 2022 ⁴⁶	2021	23.99	–	–	Mar-49	27/48	–	–	Iran (Tehran)
Joshirvani et al. 2021 ⁴⁷	May to November 2017	–	–	1.78	7.44	2.72	–	–	Iran (Hamadan)
Shahid et al. 2018 ³⁷	March and April of 2010	830	420	–	410	50	–	20	Iran (Birjand)
Farsani et al. 2018 ³⁸	October 2012 to September 2013								Iran (Ahvaz)
	autumn	2.8	–	0.19	2.85	0.39	0.3	0.61	
	normal winter	7.95	–	0.22	5.7	1.05	0.85	1.58	
	days spring	3.48	–	0.25	3.5	0.8	0.43	1.3	
	summer	2.89	–	0.2	3.1	0.66	0.35	1.1	
	autumn	4.02	–	0.51	3.67	1.55	1.45	1.79	
	Duststorm winter	8.23	–	0.87	5.78	1.8	1.58	2.35	
	spring	4.3	–	0.48	4.4	1.4	0.66	2.1	
	summer	3.45	–	0.59	3.66	1	0.72	1.5	
Di Vaio et al. 2018 ³⁹	February 2013–October 2013								Italy (Acerra)
	autumn	75.1	23.4	4.35	60.8	14.7	1.6	17.9	
	winter	76.6	17.3	4.77	50.6	19.7	3.8	16.7	
	spring	79.4	24.1	4.26	55.8	14.2	2.1	11	
Dai et al. 2015 ⁴⁸	February 2011 to January 2012	45.7	19.2	3.8	59.9	14	2.52	10.2	iron and steel industry zone background site China
		1018.6	154	4	195.2	19.2	2.4	10.7	
		794.6	117.1	3.8	158.7	7.5	1.1	6.2	

Correlation Analysis

Table 5 presents the correlation coefficients between PM₁₀ and the heavy metals. Cd exhibited the strongest correlation with PM₁₀ ($r = 0.401$, $p = 0.002$). The correlation for Zn ($r = 0.263$, $p = 0.050$) was at the borderline of statistical significance, whereas the correlation for Cu ($r = 0.246$, $p = 0.068$) was not statistically

significant. Pb, Cr, Co, and Ni exhibited positive but non-significant correlations with PM₁₀. In the correlation analysis among the metals, very strong and statistically significant relationships were observed between the metals. Cu showed the highest correlation with Pb, Cr, Co, and Ni. Pb exhibited strong correlations with Cr and Co. The correlation of Cr with Co and Ni

was also very high. Cd showed very strong and significant correlations with Zn, Cu, Pb, Cr, Co,

and Ni. These results are clearly illustrated in Figure 2.

Table 5: Pearson linear correlation coefficients for heavy metals in PM₁₀ samples.

Variables	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
(1) PM10	1.000							
(2) Zn	0.263 (0.050)	1.000						
(3) Cu	0.246 (0.068)	0.709 (< 0.001)	1.000					
(4) Cd	0.401 (0.002)	0.674 (< 0.001)	0.662 (< 0.001)	1.000				
(5) Pb	0.166 (0.221)	0.621 (< 0.001)	0.928 (< 0.001)	0.605 (< 0.001)	1.000			
(6) Cr	0.249 (0.064)	0.658 (< 0.001)	0.968 (< 0.001)	0.622 (< 0.001)	0.928 (< 0.001)	1.000		
(7) Co	0.223 (0.099)	0.516 (< 0.001)	0.879 (< 0.001)	0.563 (< 0.001)	0.849 (< 0.001)	0.953 (< 0.001)	1.000	
(8) Ni	0.249 (0.064)	0.658 (< 0.001)	0.968 (< 0.001)	0.622 (< 0.001)	0.928 (< 0.001)	1.000 (< 0.001)	0.953 (< 0.001)	1.000




Figure 2: Graphical representation of correlation patterns between PM₁₀ and heavy metals.

Comparison with International Studies

The average PM₁₀ and heavy metal concentrations in our study, along with the findings of other studies, are presented in Tables 1 and 4. The mean PM₁₀ concentration in Gorgan (69.3 $\mu\text{g}/\text{m}^3$) was lower than the values reported for Ahvaz (223.4 $\mu\text{g}/\text{m}^3$) (26), Riyadh (223.1 $\mu\text{g}/\text{m}^3$) (27), and industrial sites in Senegal (232.3 $\mu\text{g}/\text{m}^3$)

(4), but higher than those for Gijon, Spain (31 $\mu\text{g}/\text{m}^3$) (29), Lamphun, Thailand (43.0 $\mu\text{g}/\text{m}^3$) (35), and Makkah (44.5 $\mu\text{g}/\text{m}^3$) (41). The annual concentrations of Zn, Pb, and Cu were lower than the concentrations reported by Dai (2015) in China but considerably higher than those reported by Lara et al. (2023) in Spain and Rezayani et al. (2022) in Tehran. The mean concentrations of Ni

and Co were higher than the values reported in Tehran, Spain, and China (29, 44). The mean concentration of Cd was lower than the values reported for Tehran and China but higher than those reported in Spain. The mean concentration of Cr was higher than the levels reported in Spain and Tehran but lower than those in industrial regions of China.

Discussion

Variations of PM₁₀ Concentration

Results showed that PM₁₀ concentrations ranged from a minimum of 20.83 µg/m³ in early November to a maximum of 277.78 µg/m³ in early February, with a mean ± standard deviation of 69.3 ± 49.82 µg/m³ over the study period (Table 1). These values are slightly higher than the World Health Organization's 24-hour guideline of 50 µg/m³. In the study conducted by Hamidi et al. (2025)²⁵, which investigated PM₁₀ over an 11-year period (2008–2019) in Ahvaz, the mean concentration was reported as 223.4 µg/m³. Similarly, in a study by Kebe et al. (2025)⁴ in Senegal, the PM₁₀ concentrations were estimated at 232.318 and 209.854 µg/m³ at industrial and urban–coastal sites, respectively. Furthermore, Adly et al. (2025)²⁴, in a five-year study (2020–2024) in Makkah, observed an increasing trend in PM₁₀ concentrations, which rose from 127.7 µg/m³ in 2020 to 138.3 µg/m³ in 2024. Overall, the annual mean concentrations reported in all these studies are higher than those in the present investigation. Research conducted in Ahvaz and Senegal demonstrated the substantial impact of natural sources on air pollution, whereas in Makkah, the observed rise in particulate matter concentrations was primarily linked to anthropogenic activities, including population growth, traffic emissions, and construction operations. In contrast, other studies conducted by Lara et al. (2023)²⁸ in Spain, Kliengchuay et al. (2021)³⁴ in Thailand, and Munir et al. (2017)⁴⁰ in Makkah reported PM₁₀ concentrations of 31, 42.98, and 44.53 µg/m³, respectively. The observed values were not only less than those recorded in this study but also below the World Health

Organization's 24-hour guideline of 50 µg/m³. This discrepancy may reflect the varying influence of different factors, such as local meteorological conditions, emission sources, climatic parameters, policy measures, or operational programs, on PM₁₀ concentrations across different locations.

Shahsavani et al. (2025)²⁰ conducted a study in Fars province focusing on the summer season and reported a PM₁₀ concentration of 78.12 µg/m³. Elevated summer concentrations were attributed to the large amounts of suspended particles entering the atmosphere due to the desiccation of extensive areas of Bakhtegan Lake, transported by winds. In another investigation, Ahmad et al. (2023)²⁹ carried out a winter study in Pakistan and reported a PM₁₀ concentration of 279.73 µg/m³, which they linked to stagnant climatic conditions. Furthermore, Farsani et al. (2018)³⁸ examined the relationship between PM₁₀ levels on normal days and dust storm days in Ahvaz. They found that the lowest PM₁₀ concentrations on both normal and dust storm days occurred in summer (127 and 278 µg/m³, respectively), while the highest concentrations were observed in winter (189 and 742 µg/m³, respectively). Elevated PM₁₀ concentrations, particularly in winter, may be associated with the combustion of fossil fuels, coal, and wood. Conversely, lower concentrations in autumn and summer could be due to reduced human activities and the less frequent occurrence of dust storms during these seasons.

Metals Associated with PM₁₀

The mean concentrations of zinc (Zn) and lead (Pb) across all seasons exceeded the values documented in studies conducted in Fars, Tehran, Ahvaz, and Italy^{20, 43, 45, 46}. In contrast, the seasonal mean concentration of copper (Cu) was lower than the levels reported by Shahsavani et al. (2025)²⁰ and Di Vaio et al. (2018)³⁹. The mean concentration of cadmium (Cd) was also below the values reported by Di Vaio et al. (2018)³⁹ and Farsani et al. (2018)³⁸. Conversely, for cobalt (Co), the seasonal mean in the present study was higher than in these two studies. Overall, the differences reported can be attributed to variations in the

sources of these compounds and the types of sampling locations. The substantial increase in metal concentrations during winter may be due to colder conditions, increased consumption of fossil fuels, and reduced precipitation. Human activities, such as industrial operations and traffic, play a significant role in the dispersion of heavy metals. The sources of heavy metals vary; for example:

- Nickel (Ni) originates from traffic emissions, industrial activities, gas refinery, and fuel combustion⁴⁹⁻⁵¹.
- Copper (Cu) is associated with high-temperature processes, such as residential boilers and automobiles^{49, 52}.
- Lead (Pb) is primarily derived from tire wear, vehicle component corruptions, and lubricants⁵³.
- Chromium (Cr) mainly originates from agricultural activities, industrial operations, tire degradation, and greenhouse gas emissions^{54, 55}.

Consequently, heavy metals in urban air are diverse, complex, and strongly influenced by environmental conditions and seasonal variations.

Inter-Metal Correlations and Source Implications

In the correlation analysis of the metals, very strong and statistically significant relationships were observed. Positive correlations among many metals, particularly between Cu, Cr, and Ni, exhibited a very strong linear pattern with correlations approaching one. The scatter of data between PM₁₀ and Cd was more pronounced compared to other metals, indicating a significant relationship. These observations are fully consistent with the results presented in Table 4. In the study by Alharbi et al. (2024)²⁶, no significant positive correlation was observed for Cd, which differs from the observations of the current study. Similarly, the correlation results reported by Esform et al. (2022)³⁰ showed a significant relationship between PM₁₀ and Co, Cu, and Cr ($P < 0.05$), which does not align with our findings. In the Alharbi et al. (2024)²⁶ study, the correlation between Pb and Cd was significant, consistent with the present study;

however, the correlation between Cu and Cd in their study was not positive, whereas in the present study this relationship was very strong and significant. The relationship between chromium (Cr) and nickel (Ni) was very strong and significant, which is consistent with findings from the Nanjing–Shanghai highway³⁰. In the studies by Motesaddi Zarandi et al. (2019) and Esform et al. (2022), no significant correlation was observed between Pb and Cr, which does not match the results of the present study⁵⁶. Overall, the observed high positive correlations indicate significant interrelationships among the elemental constituents of PM₁₀ during the sampling period, offering a clearer perspective on their likely origins and atmospheric interactions. Furthermore, the strong correlations observed between Cu, Cr, Ni, and Pb indicate the likely presence of a common source, particularly industrial and traffic-related activities (Figure 2).

Conclusion

The findings of this study reveal that, over the study period, PM₁₀ levels in Gorgan were generally slightly above the 24-hour guideline established by the World Health Organization. Observed variations in pollutant concentrations were less influenced by seasonal changes; however, on a monthly scale, a relatively clear pattern was observed, with the highest concentrations in summer and the lowest in winter. Comparison with similar studies in other regions of the world suggests that climatic differences, local meteorological conditions, natural dust sources, and human activities can play a determining role in the levels of air pollution.

Regarding heavy metals associated with PM₁₀, the results showed that zinc (Zn), lead (Pb), and copper (Cu) exhibited the highest concentrations in the samples, typically reaching their peak levels in winter. This increasing trend can be attributed to cold weather conditions, increased fossil fuel consumption, and reduced precipitation. Additionally, strong correlations among certain metals, including Cu, Cr, Ni, and Pb, indicate the presence of common sources, particularly traffic and industrial activities.

Acknowledgments

This study is part of a research project approved by Golestan University of Medical Sciences (Project Code: 112630, IR.GOUMS.REC.1401.084). The authors express their sincere gratitude to the Vice-Chancellor for Research and Technology for their support and assistance.

Funding

This study was financially supported by Golestan University of Medical Sciences (Project Code: 112630, IR.GOUMS.REC.1401.084).

Ethical Considerations

Golestan University of Medical Sciences (IR.GOUMS.REC.1401.084).

Code of Ethics

IR.GOUMS.REC.1401.084

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and material

Not applicable.

Competing interests

The authors declare no conflicts of interest.

Author contributions

These authors contributed equally to this work and share first authorship.

This is an Open-Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt, and build upon this work for commercial use.

References

1. Kozielska B, Kaleta D. Cancer Risk Associated with Inhalation Exposure to PM₁₀-Bound PAHs and PM₁₀-Bound Heavy Metals in Polish Agglomerations. *Applied Sciences*. 2025;15(14):7903.
2. Barreira LM, Lepistö T, Salo L, et al. Comprehensive characterization of wintertime submicron aerosol in a Nordic town influenced by residential wood combustion, traffic and industrial sources. *Atmospheric Pollution Research*. 2023;14(8):101835.
3. Maceira A, Borrull F, Marcé RM. Occurrence of organic contaminants bonded to the particulate matter from outdoor air influenced by industrial activities. *Environmental Science and Pollution Research*. 2022;29(51):76644-67.
4. Kebe M, Traore A, Sow M, et al. Human health risk evaluation of particle air pollution (PM₁₀ and PM_{2.5}) and heavy metals in Dakar's two urban areas. *Asian Journal of Atmospheric Environment*. 2025;19(1):7.
5. Faraji Ghasemi F, Dobaradaran S, Saeedi R, et al. Levels and ecological and health risk assessment of PM_{2.5}-bound heavy metals in the northern part of the Persian Gulf. *Environmental Science and Pollution Research*. 2020;27(5):5305-13.
6. Kelly FJ, Fussell JC. Toxicity of airborne particles—established evidence, knowledge gaps and emerging areas of importance. *Philosophical transactions of the royal society A*. 2020;378(2183):20190322.
7. Jaafari J, Naddafi K, Yunesian M, et al. Associations between short term exposure to ambient particulate matter from dust storm and anthropogenic sources and inflammatory biomarkers in healthy young adults. *Science of The Total Environment*. 2021;761:144503.
8. Kumar RP, Prakash A, Singh R, et al. Machine learning-based prediction of hazards fine PM_{2.5} concentrations: a case study of Delhi, India. *Discover Geoscience*. 2024;2(1):34.
9. Sinha S, Kumar R, Kumar AR, et al. Elemental characteristics of respirable particulate matter in the urban atmosphere of Dehradun, Uttarakhand, India. *Curr World Environ*. 2021;16(1):319-28.
10. Zhang H, Li R, Huang C, et al. Seasonal variation of aerosol iron solubility in coarse and fine particles at an inland city in northwestern China. *Atmospheric Chemistry and Physics*. 2023;23(6):3543-59.
11. Niu S, Song X, Yu J, et al. Particle size dependent heavy metals in road dusts from Maanshan City, China. *KSCE Journal of Civil*

- Engineering. 2020;24(5):1411-23.
12. Whaley CH, Galarneau E, Makar PA, et al. How much does traffic contribute to benzene and polycyclic aromatic hydrocarbon air pollution? Results from a high-resolution North American air quality model centred on Toronto, Canada. *Atmospheric Chemistry and Physics*. 2020;20(5):2911-25.
 13. Kumar RP, Kashyap P, Kumar R, et al. Cancer and non-cancer health risk assessment associated with exposure to non-methane hydrocarbons among roadside vendors in Delhi, India. *Human and Ecological Risk Assessment: An International Journal*. 2020;26(5):1285-99.
 14. Vithanage M, Bandara PC, Novo LA, et al. Deposition of trace metals associated with atmospheric particulate matter: Environmental fate and health risk assessment. *Chemosphere*. 2022;303:135051.
 15. Swaringen BF, Gawlik E, Kamenov GD, et al. Children's exposure to environmental lead: A review of potential sources, blood levels, and methods used to reduce exposure. *Environmental research*. 2022;204:112025.
 16. Al-Lambarki M, Hasan H. Assessment of the heavy metal contents in air samples collected from the area extended between Albayda and Alquba cities (Libya). *AlQalam Journal of Medical and Applied Sciences*. 2025:695-707.
 17. Ghosh B, Padhy PK, Niyogi S, et al. A comparative study of heavy metal pollution in ambient air and the health risks assessment in industrial, urban and semi-urban areas of West Bengal, India: an evaluation of carcinogenic, non-carcinogenic, and additional lifetime cancer cases. *Environments*. 2023;10(11):190.
 18. Balali-Mood M, Naseri K, Tahergorabi Z, et al. Toxic mechanisms of five heavy metals: mercury, lead, chromium, cadmium, and arsenic. *Frontiers in pharmacology*. 2021;12:643972.
 19. Commission E. Proposal for a Directive of the European Parliament and of the Council on Ambient Air Quality and Cleaner Air for Europe. Document 52022PC0542. 2022.
 20. Shahsavani S, Shamsedini N, Mohammadpour A, et al. Air quality near Middle East's large dried lake: Heavy metal emissions, machine learning analysis, and health risks. *Physics and Chemistry of the Earth, Parts A/B/C*. 2025;137:103793.
 21. Vergara-Vásquez E, Beleño LMH, Castrillo-Borja TT, et al. Airborne particulate matter integral assessment in Magdalena department, Colombia: Patterns, health impact, and policy management. *Heliyon*. 2024;10(16).
 22. SPM PM. Sampling of ambient air for total suspended particulate matter (SPM) and PM₁₀ using high volume (HV) sampler. Center for Environmental Research Information Office of Research and Development. 1999.
 23. Aldeen SAE, Fleehe NAR. Investigating the air quality index inside two museums in Baghdad City. *Iraqi Journal of Science*. 2025.
 24. Adly HM, Saleh SA. Long-Term Trends in PM₁₀, PM_{2.5}, and Trace Elements in Ambient Air: Environmental and Health Risks from 2020 to 2024. *Atmosphere*. 2025;16(4):415.
 25. Hamidi M, Ghobadi T, Shao Y, et al. Investigation of the role of southwestern Asia dust events on urban air pollution: a case study of Ahvaz, a highly polluted city. *Scientific Reports*. 2025;15(1):21981.
 26. Alharbi HA, Rushdi AI, Bazeyad A, et al. Temporal Variations, Air Quality, Heavy Metal Concentrations, and Environmental and Health Impacts of Atmospheric PM_{2.5} and PM₁₀ in Riyadh City, Saudi Arabia. *Atmosphere*. 2024;15(12):1448.
 27. Zeb B, Ditta A, Alam K, et al. Wintertime investigation of PM₁₀ concentrations, sources, and relationship with different meteorological parameters. *Scientific Reports*. 2024;14(1):154.
 28. Lara R, van Drooge BL, Canals-Angerri A, et al. Key factors for abating particulate matter in a highly industrialized area in N Spain: fugitive emissions and secondary aerosol precursors. *Chemosphere*. 2023;341:139959.
 29. Ahmad S, Zeb B, Ditta A, et al. Morphological, mineralogical, and biochemical characteristics of particulate matter in three size fractions (PM₁₀, PM_{2.5}, and PM₁) in the urban environment. *ACS omega*. 2023;8(35):31661-74.

30. Esform A, Hassanvand MS, Naghizadeh A, et al. On the nature of heavy metals in particulate matter (PM₁₀, PM_{2.5}) and their health impact assessment for a desert city in Iran, Birjand. *Journal of Air Pollution and Health*. 2022.
31. Zeb B, Alam K, Ditta A, et al. Variation in coarse particulate matter (PM₁₀) and its characterization at multiple locations in the Semiarid Region. *Frontiers in Environmental Science*. 2022;10:843582.
32. Sirithian D, Thanatrakolsri P. Relationships between meteorological and particulate matter concentrations (PM_{2.5} and PM₁₀) during the haze period in urban and rural areas, Northern Thailand. *Air, Soil and Water Research*. 2022;15:11786221221117264.
33. Usman F, Zeb B, Alam K, et al. In-depth analysis of physicochemical properties of particulate matter (PM₁₀, PM_{2.5} and PM₁) and its characterization through FTIR, XRD and SEM–EDX Techniques in the Foothills of the Hindu Kush Region of Northern Pakistan. *Atmosphere*. 2022;13(1):124.
34. Kliengchuay W, Worakhunpiset S, Limpanont Y, et al. Influence of the meteorological conditions and some pollutants on PM₁₀ concentrations in Lamphun, Thailand. *Journal of Environmental Health Science and Engineering*. 2021;19(1):237-49.
35. Liu B, Sun X, Zhang J, et al. Characterization and spatial source apportionments of ambient PM₁₀ and PM_{2.5} during the heating period in Tian'jin, China. *Aerosol and Air Quality Research*. 2020;20(1):1-13.
36. Gupta A, Moniruzzaman M, Hande A, et al. Estimation of particulate matter (PM_{2.5}, PM₁₀) concentration and its variation over urban sites in Bangladesh. *SN Applied Sciences*. 2020;2(12):1993.
37. Shahid I, Alvi MU, Shahid MZ, et al. Source apportionment of PM₁₀ at an urban site of a south Asian mega city. *Aerosol and air quality research*. 2018;18(9):2498-509.
38. Farsani MH, Shirmardi M, Alavi N, et al. Evaluation of the relationship between PM₁₀ concentrations and heavy metals during normal and dusty days in Ahvaz, Iran. *Aeolian Research*. 2018;33:12-22.
39. Di Vaio P, Magli E, Caliendo G, et al. Heavy metals size distribution in PM₁₀ and environmental-sanitary risk analysis in Acerra (Italy). *Atmosphere*. 2018;9(2):58.
40. Munir S, Habeebullah TM, Mohammed AM, et al. Analysing PM_{2.5} and its association with PM₁₀ and meteorology in the arid climate of Makkah, Saudi Arabia. *Aerosol and Air Quality Research*. 2017;17(2):453-64.
41. Alam K, Azam M, Shahid I, et al. Source apportionment and characterization of particulate matter (PM₁₀) in urban environment of Lahore. *Aerosol and Air Quality Research*. 2014;14(7):1851-61.
42. Sharma S, Mandal T, Saxena M, et al. Source apportionment of PM₁₀ by using positive matrix factorization at an urban site of Delhi, India. *Urban climate*. 2014;10:656-70.
43. karimi sr. The amount of suspended particles containing heavy metals in the air of the 21st district of Tehran, 2021 autumn. *Iranian Journal of Biological Sciences*. 2023;4(17):61-76.
44. Rezayani N, Mir Mohammadi M, Mehrdadi N. Determination of selected heavy metals in air samples and human health risk assessment in Tehran city, Iran. *Iran J Chem Chem Eng Research Article Vol*. 2022;41(8).
45. Karimi SR, Mansouri N, Taghavi L, et al. Investigation of Heavy Metal Concentrations in Airborne Particles in District 21 of Tehran. *Journal of Research in Environmental Health*. 2022;8(2):136-47.
46. Karimi SR, Mansouri N, Taghavi L, et al. The receptor of heavy metals in total particulate matter with UNMIX determine the contribution model in 21st district of Tehran. 2022.
47. Joshirvani A, Samarghandi MR, Leili M. PM₁₀ Concentration, Its Potentially Toxic Metals Content, and Human Health Risk Assessment in Hamadan, Iran. *CLEAN–Soil, Air, Water*. 2021;49(6):2000174.
48. Dai Q-L, Bi X-H, Wu J-H, et al. Characterization and source identification of heavy metals in ambient PM₁₀ and PM_{2.5} in an

- integrated iron and steel industry zone compared with a background site. *Aerosol and Air Quality Research*. 2015;15(3):875-87.
49. Bi C, Chen Y, Zhao Z, et al. Characteristics, sources and health risks of toxic species (PCDD/Fs, PAHs and heavy metals) in PM_{2.5} during fall and winter in an industrial area. *Chemosphere*. 2020;238:124620.
50. Jiang N, Liu X, Wang S, et al. Pollution characterization, source identification, and health risks of atmospheric-particle-bound heavy metals in PM₁₀ and PM_{2.5} at multiple sites in an emerging megacity in the central region of China. *Aerosol and Air Quality Research*. 2019;19(2):247-71.
51. Nargis A, Habib A, Islam MN, et al. Source identification, contamination status and health risk assessment of heavy metals from road dusts in Dhaka, Bangladesh. *Journal of Environmental Sciences*. 2022;121:159-74.
52. Kamani H, Hoseini M, Safari GH, et al. Study of trace elements in wet atmospheric precipitation in Tehran, Iran. *Environmental monitoring and assessment*. 2014;186(8):5059-67.
53. Hou S, Zheng N, Tang L, et al. Pollution characteristics, sources, and health risk assessment of human exposure to Cu, Zn, Cd and Pb pollution in urban street dust across China between 2009 and 2018. *Environment international*. 2019;128:430-7.
54. Jiang Y, Ma J, Ruan X, et al. Compound health risk assessment of cumulative heavy metal exposure: a case study of a village near a battery factory in Henan Province, China. *Environmental Science: Processes & Impacts*. 2020;22(6):1408-22.
55. Cheng X, Huang Y, Zhang S-P, et al. Characteristics, sources, and health risk assessment of trace elements in PM₁₀ at an urban site in Chengdu, Southwest China. *Aerosol and Air Quality Research*. 2018;18(2):357-70.
56. Motesaddi Zarandi S, Shahsavani A, Khodagholi F, et al. Concentration, sources and human health risk of heavy metals and polycyclic aromatic hydrocarbons bound PM_{2.5} ambient air, Tehran, Iran. *Environmental geochemistry and health*. 2019;41(3):1473-87.