



## Quantum Dots and Their Application in Water and Wastewater Treatment

Saeid Ahmadzadeh <sup>1\*</sup>, Maryam Dolatabadi <sup>2</sup>, Roya Malekhamadi <sup>2</sup>, Ali Rasooli <sup>2</sup>

<sup>1</sup> Neuroscience Research Center, Institute of Neuropharmacology, Kerman University of Medical Sciences, Kerman, Iran.

<sup>2</sup> Environmental Sciences and Technology Research Center, Department of Environmental Health Engineering, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.

### ARTICLE INFO

#### LETTER TO EDITOR

#### Article History:

Received: 10 January 2023

Accepted: 21 February 2023

#### \*Corresponding Author:

Saeid Ahmadzadeh

#### Email:

chem\_ahmadzadeh@yahoo.com

#### Tel:

+98 34 31325241

**Citation:** Ahmadzadeh S, Dolatabadi M, Malekhamadi R, et al. *Quantum Dots and Their Application in Water and Wastewater Treatment*. J Environ Health Sustain Dev. 2023; 8(1): 1862-4.

In recent years, the use of nanoparticles has grown tremendously in various fields, including medicine, industry, agriculture, mechanics, and water and wastewater treatment. As a functional nanoparticle, Quantum dots (QDs), have recently been introduced to the field of water and wastewater treatment <sup>1</sup>. QDs comprising of 200 to 10,000 atoms have demonstrated great versatility due to their small diameter of 2 to 10 nm, and function as a semi-conductor nanocrystal by emitting light after excitation. The QDs characteristics are controlled by changing the molecular structure according to the individual's desirable usage <sup>2</sup>. The size, shape, and structure of QDs control their electrical properties and peak emission frequency <sup>3,4</sup>.

The electrical conductivity of QDs show significant changes by applying external stimuli, such as an electric field or light radiation. QDs are commonly categorized into two groups; metal quantum dots (MQDs); and carbon quantum dots (CQDs). MQDs such as PbS, ZnS, InP, InAs, CdTe, CdSe, and CdS emit light after exciting the electrons in accordance with a certain wavelength of light by an external source. CQDs with a size of less than 10

nm and discrete quasi-spherical shape have been considered as an attractive group of nanoparticles <sup>5,6</sup>.

Graphene QDs, carbon nanodots, and polymer dots are categorized as CQDs and played a valuable and substantial role in aqueous mediums treatment, due to their attractive properties of low toxicity, fluorescence activity, eco-friendliness, low cost, unique optical properties, and chemical stability <sup>7,8</sup>. The high fluorescence properties of CQDs have made them suitable for various applications, and they have received considerable attention from researchers as fluorescent carbon dots. Various approaches have been developed for the preparation of CQDs, including hydrothermal methods, radiation and ultrasound procedures, electrochemical strategies, and microwave techniques. Hydrothermal techniques are most commonly employed for the synthesis of CQDs, and more progressive techniques have been recently developed for their preparation <sup>9,10</sup>.

CQDs have recently been employed as an efficient adsorbent for removal of organic pollutants such as dyes <sup>11,12</sup>, and inorganic contaminants such as heavy metals, pharmaceutical materials <sup>13</sup>, p-nitrophenol <sup>14,15</sup>, pesticides <sup>1,16</sup>, and pathogens <sup>17,18</sup>.

They have been also used as an active catalyst for photo-catalytic treatment processes of water and wastewater<sup>6,8</sup>.

Although CQDs have been extensively studied by researchers, there are some features and challenges in their utilization and usage in water and wastewater treatment processes which need to be resolved. These include a comprehensive understanding of CQDs performance, cost effectiveness and simple synthesis approaches for preparation of CQDs with green carbon precursors, and regeneration investigations.

This is an Open-Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt, and build upon this work for commercial use.

## References

1. Ghaffarkhah A, Hosseini E, Kamkar M, et al. Synthesis, applications, and prospects of graphene quantum dots: A comprehensive review. *Small*. 2022;18(2):2102683.
2. Bai Y, Hao M, Ding S, et al. Surface chemistry engineering of perovskite quantum dots: strategies, applications, and perspectives. *Mater Adv*. 2022;34(4):2105958.
3. Das R, Bandyopadhyay R, Pramanik PJM. Carbon quantum dots from natural resource: A review. *Mater Today Chem*. 2018;8:96-109.
4. Xu Q, Niu Y, Li J, et al. Recent progress of quantum dots for energy storage applications. *Carbon Neutrality*. 2022;1(1):13.
5. Rao VN, Reddy NL, Kumari MM, et al. Sustainable hydrogen production for the greener environment by quantum dots-based efficient photocatalysts: a review. *J Environ Manage*. 2019;248:109246.
6. Heng ZW, Chong WC, Pang YL, et al. An overview of the recent advances of carbon quantum dots/metal oxides in the application of heterogeneous photocatalysis in photodegradation of pollutants towards visible-light and solar energy exploitation. *J Environ Chem Eng*. 2021;9(3):105199.
7. Wang L, Jiang H, Wang H, et al. MXenes as Heterogeneous Fenton-like Catalysts for Removal of Organic Pollutants: A Review. *J Environ Chem Eng*. 2022;10(6):108954.
8. Wang L, Luo D, Yang J, et al. Metal-organic frameworks-derived catalysts for contaminant degradation in persulfate-based advanced oxidation processes. *J Clean Prod*. 2022;375(15):134118.
9. Tian L, Li Z, Wang P, et al. Carbon quantum dots for advanced electrocatalysis. *J Energy Chem*. 2021;55:279-94.
10. Devi P, Saini S, Kim K, et al. The advanced role of carbon quantum dots in nanomedical applications. *Biosens Bioelectron*. 2019;141:111158.
11. Hui KC, Ang WL, Sambudi NS, et al. Nitrogen and bismuth-doped rice husk-derived carbon quantum dots for dye degradation and heavy metal removal. *J Photochem Photobiol*. 2021;418:113411.
12. Kang Z, Jia X, Zhang Y, et al. A review on application of biochar in the removal of pharmaceutical pollutants through adsorption and persulfate-based AOPs. *Sustainability*. 2022;14(16):10128.
13. Li C, Zhao Z, Fu S, et al. Polyvinylpyrrolidone in the one-step synthesis of carbon quantum dots anchored hollow microsphere Bi<sub>2</sub>WO<sub>6</sub> enhances the simultaneous photocatalytic removal of tetracycline and Cr (VI). *Sep Purif Technol*. 2021;270:118844.
14. Fan P, Zhang X, Deng H, et al. Enhanced reduction of p-nitrophenol by zerovalent iron modified with carbon quantum dots. *Appl Catal*. 2021;285:119829.
15. Yoosefian M, Sabaei S, Etminan N. Encapsulation efficiency of single-walled carbon nanotube for Ifosfamide anti-cancer drug. *Comput Biol Med*. 2019;114:103433.
16. Huang R, Yang J, Cao Y, et al. Peroxymonosulfate catalytic degradation of persistent organic pollutants by engineered catalyst of self-doped iron/carbon nanocomposite derived from waste toner powder. *Sep Purif Technol*. 2022;291:120963.

17. Anand KV, Subala AS, Sumathi KS, et al. A review on the removal of dye, pesticide and pathogens from waste water using quantum dots. *European Journal of Advanced Chemistry Research*. 2020;1(5):1404.
18. Mirhaji E, Afshar M, Rezvani S, et al. Boron nitride nanotubes as a nanotransporter for anti-cancer docetaxel drug in water/ethanol solution. *J Mol Liq*. 2018;271:151-6.