

Journal of Environmental Health and Sustainable Development



Indoor and Outdoor Air Fungus Bioaerosols in Khorramabad Day Care Child Centers Western of Iran, 2018

Asghar Sepahvand¹, Katayan Salim^{2*}, Edris Hoseinzadeh³, Khadijeh Jafari^{4,5}, Rezvan Mohammadrezaei Khorramabadi⁶

ARTICLE INFO

ORIGINAL ARTICLE

Article History:

Received: 28 May 2020 Accepted: 10 July 2020

*Corresponding Author:

Katayan Salim

Email:

salmkate@yahoo.com

Tel:

+989163673104

Keywords:

Fungi, Air Pollution, Indoor, Child Day Care Centers, Iran.

ABSTRACT

Introduction: the aim of this study was to determine the quantity and quality of indoor and outdoor air fungus bioaerosols in Khorramabad day care child centers

Materials and Methods: A total of 180 air samples were collected from 10 centers in 2018. The samples included 7 indoor and 2 outdoor sampling points. The total number of children was 580. Sampling of fungal bioaerosols was performed by the ZEFON pump (ZEFON factory, USA) with a flow rate of 28.3 L/min. The Sabouraud Dextrose Agar containing chloramphenicol was used as the culture medium. Relative humidity and temperature were measured by a Hygro-Thermometer (TES-1360A- Taiwan-made Humidity and temperature meter).

Results: The results showed that 96.1 % of the samples were positive and had grown colonies. The highest amount of fungal agents in the indoor air and outdoor air were 175.58 CFU/m³ and 274.56 CFU/m³ in May, while the lowest rates were 3.4 CFU/m³ and 7.8 CFU/m³ in July, respectively. *Aspergillus niger* and *Mucor* were the most highly abundant fungus genera, while *Fusarium* was the lowest one. In all samples, the I/O (indoor/outdoor) ratio was more than 1; so, fungal bioaerosols in indoor environments were dominant than the outdoor fungal bioaerosols. The relationship of fungal bioaerosols with RH and T(°C) was significant (P-value = 0.001).

Conclusion: Generally, the amount of contamination is considerable in the studied day care child centers. Therefore, ventilation modification is recommended by a purifier filter. Moreover, the ventilation conditions and favorable air standards should be monitored continuously by supervisory authorities.

Citation: Sepahvand A, Salim K, Hoseinzadeh E, et al. *Indoor and Outdoor Air Fungus Bioaerosols in Khorramabad Day Care Child Centers Western of Iran*, 2018. J Environ Health Sustain Dev. 2020; 5(3): 1077-90.

Introduction

Bioaerosols are a mixture of small particles including dust, microbes, fragments, and their fragments residues. The pathology effects of these bioaerosols depend on the size, concentration, physicochemical properties, and distribution of their particles ¹ since the micro- to nano-scale size bioaerosols penetrate the lungs and circulation

¹ Razi Herbal Medicines Research Center, Lorestan University of Medical Sciences, Khorramabad, Iran.

² Department of Environmental Engineering, Ahvaz Branch, Islamic Azad University, Ahvaz, Iran.

³ Student Research Committee, Saveh University of Medical Sciences, Saveh, Iran.

⁴ Environmental Science and Technologhy Research Center, Department of Environmental Health Engineering, School of Public Health, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.

⁵ Environmental Health Engineering Officer, Karafarinan Ava Salamat Compony, Shiraz University Of Medical Sciences, Qaemiyeh, Iran.

⁶ Department of Nursing, School of Nursing and Midwifery, Lorestan University of Medical Sciences, Khorramabad, Iran.

[DOR: 20.1001.1.24766267.2020.5.3.7.6]

DOI: 10.18502/jehsd.v5i3.4279

system easily in different parts of the body ². Thus, the type, source, density, and complication effects of bioaerosols should be studied on the personal health ³. Bioaerosol production was documented by human activities including sneezing/coughing, room floor washing, cleaning toilets and makeup tables, walking/talking, etc. ⁴.

Indoor air pollution leads to a variety of adverse effects health including acute respiratory infections, middle ear infections, gastro-intestinal diseases, skin problems, invasive bacterial infections, eye diseases, throat inflammation, skin and nose problems, dizziness, asthma, and allergic reactions ⁵. Such health effects bring about a significant economic burden on the involved countries. According to the studies of bioaerosol pollution, indoor air is affected by outdoor air pollution since people are usually in closed environment for a long time ⁶.

After investigation of the concentration and size of airborne bioaerosols, researchers reported that climatic conditions such as air temperature, relative humidity, velocity wind, sandstorms, as well as proximity to the sea, drought, and seasons influence the growth and increase bioaerosol concentration ⁷. Moreover, the relationship among fungal concentration, suspended particles, and meteorological parameter was determined ⁸.

Fungal microorganisms are the most wellknown and diverse species in indoor and outdoor environments. The concentration and type of airborne micro-fungus in the indoor and outdoor environment depend on biological environmental factors 9, 10. Fungal colonies may release single spores, cluster of spores, and small fungal fragments in the environment 11. Some studies showed that fungi had the highest percentage of isolated bioaerosols that produced several types of mycotoxins and volatile organic compounds (VOCs) 11.

Children are considered a risk group because they are more vulnerable to pollutants than adults ¹². Acute respiratory infections and ear infections are most common among preschool children who were kept in kindergartens rather than children who were in home ⁶. One of the reasons for sensitivity of the children's age group is that inhalation of air per weight unit in resting infants is twice higher than that of an adult, which is due to the lack of development in children's immune system and lungs 12-15. Children also have higher heart rates than adults because they allow the absorbed substances in the blood to permeate the tissues rapidly ¹⁶.

Some studies investigated the amount of airborne fungi in indoor air of the child day care centers (CDCC) 17. For example, Hoseinzadeh et al. studied the airborne fungal flora in the indoor air of DCCs in Delfan. They showed that the most isolated fungal species was Penicillium Spp. The maximum and minimum of total fungi (CFU) were 372 CFU or 1314 CFU/m³ and 91 CFU or 322 CFU/m³, respectively. However, they just studied indoor air not the outdoor air. Evidence suggests that indoor air pollution can lead to respiratory damages; inflammation in the skin, eyes, lungs, throat, and upper respiratory tract problems, dyspnea, dizziness, and allergic reactions ¹⁸. Typical indoor environments where children are kept for a long time include schools, DCCs, and homes ¹⁹. Children under the age of 6 years are in the kindergartens at least 8 hours a day and are divided into several age groups: infants (less than 1 year old), toddlers (1-3 years old), and preschool children (3-5 years old) ⁶.

Children kept in DCCs are often exposed to respiratory infections for a long time. Such respiratory infections are related to the problems caused by fungal molds in the building 20, 21, which are called the sick building syndrome 20-22. Therefore, childhood atopic diseases can occur due to indoor allergens such as tiny worms and garbage, fungi, beetles, and pets such as dogs and cats 23. In Iran, children spend a lot of time in DCCs. Therefore, it is important to pay attention to bioaerosols due to their risk for children ⁶.

Numerous studies were conducted on the microbial quality monitoring in indoor and outdoor environments in the developed and developing countries. However, limited studies were conducted on the fungal contamination of the children kept in DCCs in Iran. Given the necessity

to ensure the health of children kept in DCCs and to improve the health status of these places, fungal bioaerosols were investigated in the indoor and outdoor air in children DCCs of Khorramabad.

Materials and methods

Sample sites description

This study was conducted in Khorramabad City, the capital of Lorestan province in Iran. The population of this city was 506471 people in 2016, according to the latest census of Iran. Khorramabad City is located at the 33° 48′ N, 48° 35′ E ^{24, 25}. It has a mild and humid climate. The average annual precipitation is 509 mm. In terms of geomorphology, the study area is mountainous with many mountains around the city of Khorramabad. The average daily temperature in Khorramabad City is 17.2 °C and the speed of the west wind is 10 km/h with an RH of 14%.

The method and sampling areas

Khorramabad City was divided into 4 regions including north, west, south, and center according to the DCCs density, population, and sources of air pollution. Due to the objectives of the study and sampling method, the traffic condition of the city was also considered. In this regard, the required information available at the traffic police office of Khorramabad City was divided into three sections in terms of traffic including high, medium, and low traffic areas. In this study, the total population of children in DCCs was 580 including 272 females and 308 males (M = 308 and F = 272).

From 215 DCCs in Khorramabad, 10 DCCs were selected randomly in several stages. To consider the effect of traffic on air pollution, 5 DCCs were selected from high traffic areas and 5 DCCs were selected from the low traffic areas in the north, south, west, and east regions of Khorramabad. According to the statistical calculations, 180 samples (from indoor and outdoor environments of the children DCCs) were collected during the two seasons of spring and

summer in 2018. The samples were taken in the morning (from 9 to 11 AM) with doors and windows closed for about 2 hours and without presence of children in the sampling places. The studied variables included airborne fungus concentrations, type of fungi, age of the child, gender, type of the children DCCs, relative humidity, temperature, season of the year, children DCCs location, sampling time, and wind speed.

The DCCs' sampling sites included seven indoor air samples and 2 stations considered as the outdoor sampling sites of the DCCs. Demographic data of the DCCs including the DCCs' type, address, number of children, and number of teachers, children's age and gender, sampling time, month, and period, as well as geographical location were collected through in-person interviews by questionnaires.

The prepared samples were transferred to Razi Herbal Medicine Research Center of Lorestan University of Medical Sciences for diagnosis and then the species and density of the fungal bioaerosols were determined. To determine the relationship between the colony count environmental conditions, we determined the concentration of the suspended particles and measured the Τ, RH, and wind speed simultaneously through fungal sampling. The weather data were also collected from the Khorramabad Meteorological Organization. **Figure** geographical location of Khorramabad City in Lorestan province depicted.

Children under the age of 6 years, who spend at least 8 hours a day in DCCs, were divided into several age groups including: infants (less than one year old), toddlers (1-3 years old), and preschool children (3-5 years old). In the studied DCCS, air conditioners as well as natural ventilation by windows (often open) were used in warm seasons of the year. In the cold seasons, gas heaters were used.

10/9

1080




Figure 1: Geographical location of the studied DCCs in Khorramabad City

Sampling fungal agents

The study method was adopted from the literature ^{6, 11, 26}, where active air sampling method and single stage Anderson method were applied. The sampling method included collision of bioaerosols with single stage Andersen's impactor (Quick Take-30, USA). The volume of air required in this study was based on the NIOSH0800 standard and the Andersen's impactor 1ft³/min method, which was 28.3 l/min. Each sampling lasted 2.5 min in this study, but a shorter sampling time is necessary in heavily contaminated areas. Later, 10cm plates containing a culture medium of Sabouraud dextrose agar (SDA) with 0.5 g/l chloramphenicol (made in Spain, growth medium type) were placed in Andersen's impactor (Quick Take-30, USA) according to the instructions provided by the manufacturer company. device was placed at the height of the breathing zone of children, i.e., 0.6-1.2 m. Sampling was performed in all days of the week and in different random places and times 27. Corn meal agar Germany) was used to stimulate (merck,

sporulation of the fungi. Zefon sampling pump (manufactured by SKC in USA) was used with a maximom sampling capability of 30 l/min for microbial sampling. Furthermore, the Andersen's impactor was used to make suction (Quick Take-30, USA). Digital TES-1360 (Germany) was applied for measuring %RH and T (°C). Before starting the measurement operation, the TES-1360 was calibrated by the licensed company.

Calibration of the Andersen's impactor Flow **Impedance** was performed through Rotameters at a flow rate of 28.3 l/min before each sampling. The selection of sampling sites was based on the role and importance of sampling point conditions in indoor and outdoor fungal contamination. All the necessary tools and equipment were autoclaved for 15 min under standard temperature and sterilized by alcohol 70% isopropanol before sampling. After sampling, to prevent the secondary contamination and error, we closed the plates' lids and transferred them into sterile packages. Later, the plates were incubated at temperature of 25-27°C (laboratory temperature

was room temperature) for 3-7 days ²⁸⁻³³. After the growth and appearance of fungi on the surface of the culture medium, fungi were identified based on macroscopic and microscopic characteristics ³⁴.

To calculate the number of fungal spores per cubic meter, the following formula was used with the quantum value defined as 1 CFU/m^3 35, 36.

Eq 1. CFU m⁻³ =
$$\frac{T \times 1000}{t \times F(\frac{L}{min})}$$

T: The number of colonies on the surface of fungal culture medium (SDA)

1000: Conversion coefficient of flow (l/m³)

t: Sampling time (min)

L: Sampling pump flow (l/min)

Air temperature, relative humidity (Testoterm, Germany), and wind speed (Utron, AM-4201) were performed every 3 min once in the sampling process. In the case that the difference between T and pressure used for calibration was significant, sampling was modified using the ideal gas law.

Macroscopic and microscopic properties

Different macroscopic characteristics of the colonies grown on the Sabouraud dextrose agar were investigated with chloramphenicol: colonies' state and shape, colonial surface view, colony consistency, colony count, colony color, back and front color of the colony, tissue type, exudative secretion droplets, and sclerotia. The microscopic checkup of the fungal species was also carried out using crushing colony methods and fungus culture on lam, using Lactophenol Coton Blu (LPCB) solution. Furthermore, slide cultures and needle mounts (tease mounts) were used for microscopic determination according to Aneja K et al.³⁴. To observe the non-sexual spore formation, quiddity arrangement, spore manufacturer cells, as well as growth in different temperatures were considered ²⁷. When the number of colonies was high and difficult to count, **back**-calculated image, texture, pigments, and topography of colony, as well as exudate and sclerotium of the plates were used. To determine the fungal genera and species, methods applied in the related literature were used^{34, 37-39}.

Statistical analysis

Control samples were provided to ensure the accuracy of the results. At first, the normality of data assessed by the Kolmogorov-Smirnov and the Shapiro-Wilk tests. If the data were not normal (Sig. value < 0.05), Spearman and Kruskal–Wallis nonparametric tests were applied to analyze the comparison between variables and DCCs, respectively. The Pearson test applied when the data were normal. All statistical analyses were conducted using SPSS software (version 21) at significant level of 0.05.

Comparisons were made using the guidelines and reference standards for RH and T. According to the American Society of Heating, Refrigerating, and Air-Conditioning Engineers (ASHRAE) standard, T ranges 20-29 °C in the winter and 22.8-26.8 °C in the summer. Furthermore, RH was within the range of 60-30 percent ⁴⁰. Charts were designed using EXCELL 2016.

Results

Comfort parameters

The mean temperature was 32.73 ± 7.02 °C and the mean RH was 33.19 ± 2.66 in the studied DCCs. According to the sampling site, the highest T °C was in the summer with an average of 38.80 ± 3.99 . The mean RH content in this study was 33.19 ± 2.66 , which was generally in accordance with the ASHRAE standard. Table 1 shows the RH content, T rate, and volume of colonies based on the season and geographical location of DCCs.

Table 1: Average temperature (°C), RH (%), and number of colonies per unit volume (CFU/m³) in the geographical location of DCCs in the Khorramabad City

Geographical location	T (°C)	% RH	Volume (CFU/m ³)
North	2.94 ± 26.6	1.93 ± 34.93	111.92 ± 201.78
South	3.99 ± 38.80	2.08 ± 31.44	100.02 ± 187.65
Center	2.94 ± 26.67	1.93 ± 34.93	111.92 ± 201.78
West	3.99 ± 38.80	2.08 ± 31.44	100.02 ± 187.65
Total	7.02 ± 32.73	2.66 ± 33.19	106.08 ± 194.71

Fungal bioaerosols

Table 2 shows the distribution of fungal species type in indoor and outdoor DCCs based on the research variables. Aspergillus niger was the most frequent species in the samples, which was followed by Mucor, Penicillium, Aspergillus flavus, Rhizopus, Cladosporium, Alternaria, and Fusarium species, respectively.

According to the geographical location of DCCs, south and west areas were the warmest

location with the lowest average of RH (%). The northern and central regions had the lowest temperature and highest RH (%) average content.

Furthermore, the Chi-Square test showed no significant difference between different geographical locations and the type of fungal contamination, apart from Aspergillus flavus fungus, which was found in the western regions of the city.

Table 2: Distribution of the selected samples from DCCs in Khorramabad based on the contamination with fungal agents (CFU/m³), season, type of DCC, geographical location, sampling point, and sample selection space

Contamination (CFU/m³)		Without colonies (CFU/m³)		Under 15 colonies (CFU/m³)		15-30 colonies (CFU/m³)		Higher than 30 colonies (CFU/m³)		Total		P- value
	Season	Number	Percent	Number	,	Number	Percent	,	,	Number	Percent	
Season	Spring	5	(5.6)	35	(38.9)	41	(45.6)	9	(10)	90	(100)	0.292
	Summer	2	(2.2)	41	(45.6)	43	(47.8)	4	(4.4)	90	(100)	
Type of DCC	Governmental	3	(3.3)	31	(34.4)	46	(51.1)	10	(11.1)	90	(100)	0.064
	Personal	4	(4.4)	45	(50)	38	(42.2)	3	(3.3)	90	(100)	
	North	0	(0)	28	(53.8)	21	(40.4)	3	(5.8)	52	(100)	0.092
ohica ion	South	2	(3.8)	15	(28.3)	29	(54.7)	7	(13.2)	53	(100)	
Geographical location	Center	2	(5.4)	15	(40.5)	19	(51.4)	1	(2.7)	37	(100)	
Ge	West	3	(8.8)	17	(50)	12	(35.3)	2	(5.9)	34	(100)	
	Hall	0	(0)	9	(45)	10	(50)	1	(5)	20	(100)	0.356
	corridor	1	(5)	9	(45)	10	(50)	0	(0)	20	(100)	
	W.C	2	(10)	9	(45)	9	(45)	0	(0)	20	(100)	
ıts	Kitchen	0	(0)	11	(55)	8	(40)	1	(5)	20	(100)	
l poir	Shahid Nobaveh 1	2	(10)	7	(35)	10	(50)	1	(5)	20	(100)	
Sampled points	Shahid Nobaveh 2	2	(10)	7	(35)	7	(35)	4	(20)	20	(100)	
	Shahid Nobayeh 3	0	(0)	8	(40)	8	(40)	4	(20)	20	(100)	
	Outdoor 1	0	(0)	7	(35)	12	(60)	1	(5)	20	(100)	
	Outdoor 2	0	(0)	9	(45)	10	(50)	1	(5)	20	(100)	
led es	Indoor	7	(5)	60	(42.9)	62	(44.3)	11	(7.9)	140	(100) (100)	
Sampled places	Outdoor	0	(0)	16	(40)	22	(55)	2	(5)	40	(100)	0.356

A significant relationship was found between the frequency of fungal bioaerosols and temperature (P-value < 0.01).

Based on the findings, 6% of the samples were contaminated with Fusarium fungi, which was only observed in spring (P-value = 0.014); but contamination with other fungal agents did not show a significant relationship with the season.

In spring, only 5.6% of the samples had no contamination; 94.4% of the samples had less than 15 to more than 30 fungus colonies. This percentage dropped to 2.2 in the summer. However, no significant difference was observed between the two seasons in terms of contamination with fungal bioaerosols.

No significant difference was found between governmental and private DCCs. In the north region, all samples were contaminated, but 5.4%, 8.8%, and 3.8% of DCCs were without contamination in the center, west, and south regions of the city, respectively. However, the observed differences were not statistically significant (P-value = 0.092). No significant difference was found between the samples taken from the indoor and outdoor areas and in various DCCs sites in terms of contamination with fungal

agents. The highest contamination agent was *Aspergillus niger* (91.1%) and the lowest was 3.8% for *Fusarium*.

The amount of fungal agents was investigated for indoor or outdoor spaces and sampling seasons. The highest amount of fungal agents in the indoor environment was observed at 175.58 CFU/m³ in May, while its lowest value was 3.4 CFU/m³ in July. The highest amount of fungal agents in the outdoor environment was measured at 274.56 CFU/m³ in May and its lowest value was 7.8 CFU/m³ in July. In this study, no significant difference was found between the sampling sites such as hall, shahid Nobaveh three classrooms, WC, kitchen, corridor, and outdoor stations (P-value = 0.356).

Comparison of the selected samples from outdoor and indoor spaces indicated that the outdoor samples were significantly more contaminated with *Rhizopus* fungus than indoor samples (P-value = 0.001). No relationship was found between contamination with other fungal agents and open or closed spaces, except for *Rhizopus* (Table 3). In all samples, the I/O (indoor air/outdoor air) ratio was greater than 1.

Table 3: Frequency distribution of the observed fungi in samples selected from DCCs in Khorramabad City based on fungus type and sampling space

Sampling spaces Type of fungi		Outdoor		Indoor		Total		D l
		Number	Percent	Number	Percent	Number	Percent	P-value
Aspergillus niger	Positive	34	(20.7)	130	(79.3)	164	(100)	0.124
	Negative	6	(37.5)	10	(62.5)	16	(100)	
Aspergillus flavus	Positive	9	(23.1)	30	(76.9)	39	(100)	0.885
	Negative	31	(22)	110	(78)	141	(100)	
Penicillium	Positive	17	(30.4)	39	(69.6)	56	(100)	0.078
	Negative	23	(18.5)	101	(81.5)	124	(100)	
Cladosporium	Positive	4	(30.8)	9	(69.2)	13	(100)	0.442
	Negative	36	(21.6)	131	(78.4)	167	(100)	
Mucor	Positive	24	(21.6)	87	(78.4)	111	(100)	0.806
	Negative	16	(23.2)	53	(76.8)	69	(100)	
Fusarium	Positive	2	(33.3)	4	(66.7)	6	(100)	0.506
	Negative	38	(21.8)	136	(78.2)	174	(100)	
Alternaria	Positive	3	(37.5)	5	(62.5)	8	(100)	0.288
	Negative	37	(21.5)	135	(78.5)	172	(100)	
D1.:	Positive	11	(50)	11	(50)	22	(100)	0.001
Rhizopus	Negative	29	(18.4)	129	(81.6)	158	(100)	0.001

108.

Discussion

The highest T ($^{\circ}$ C)was in the summer with an average of 38.80 \pm 3.99. The average temperature in this study was contrary to the ASHRAE standard 6 .

A number of studies pointed out the relationship between relative humidity and bioaerosols in indoor air 41. A direct relationship was found agents between fungal and meteorological parameters including relative humidity and temperature in indoor environment as well as temperature, relative humidity, and wind speed in the outdoor environment. This is probably due to ventilation and indoor air pressure as well as closed doors of the DCCs. In addition, the presence of Aspergillus niger may be due to the suitable condition of DCCs to the growth of Aspergillus niger.

The mean RH content in this study was 33.19 ± 2.66 , which was generally in accordance with the ASHRAE standard. The approximate RH of 20 to 60% leads to comfort for the human body. Low or high RH levels can lead to physical problems, an RH content of less than 20% causes severe ocular stimulation, and high RH levels can reduce the severity of asthma ^{6, 42}. The world health organization recommends that people should live in moderately humid environments to prevent the occurrence of respiratory infections associated with low or high relative humidity ²³.

All isolated fungi require an average rate of RH to grow and reproduce. The lowest water activity was in primary colonizers such as *Penicillium* spp., *Alternaria*, and the tertiary colonizers (only in the *Fusarium*). Based on fungal water requirements, most isolated fungi such as *Cladosporium*, *Aspergillus flavus*, *Aspergillus niger*, *Rhizopus*, and *Mucor* can be divided into secondary colonies that require water activity of about 0.8-0.9% ⁴³.

Most fungi isolated in this study are important allergens that can cause symptoms, such as sick building syndrome, respiratory diseases, and childhood systemic infections ^{20, 21}.

According to the geographical location of DCCs, the ones located in the south and west areas were the warmest location with the lowest average

of RH (%). The northern and central regions had the lowest average temperature and highest RH (%) content. The Chi-Square test showed no significant difference between different geographical locations and the type of fungal contamination, apart from Aspergillus flavus fungus found in the western regions of the city. In another study in São Paulo city of Brazil (2017), Aspergillus flavus, as an important source of allergens, was abundant 44. Park Donguk et al. investigated airborne microbial and fungal contamination in 70 classes in 17 DCCs. They showed that the fungus concentration was greater than 1000 CFU/m³. According to the multivariate regression tests, fungus sampling and ventilation efficiency can affect the microbial concentration in the samples ⁴⁵.

A significant relationship was observed between the frequency of fungal bioaerosols and temperature (P-value < 0.01). Therefore, environment temperature is one of the important factors in determining the type of fungal species 6 . *Fusarium* fungi contaminated 6% of the samples and was only observed in spring (P-value = 0.014), which can be due to the sampling location that was near the corn fields near the Khorramabad city 46 . Contamination with other fungal agents did not have a significant relationship with the season.

In the study by Hoseini et al., the maximum concentration of airborne fungi in the outdoor air was $275 \text{ CFU/m}^3 \text{ (minimum} = 21 \text{ CFU/m}^3 \text{) for}$ Imam Khomeini station that is similar to the result of the present study. Furthermore, the minimum concentration was 127 CFU/m³ in office area as an indoor environment (maximum = 1060 CFU/m³), which may be due to the place of sampling (subway stations) 31. Hoseinzadeh et al. invetsigated the hospitals in Hamedan and showed that the total mean concentration of fungal bioaerosols was 12.56 CFU/m³. This rate is lower than 100 CFU/m ³ set as a guideline by ACGIH/Guidelines for assessing bioaerosols in 1989 (according to consensus) and less than 50 CFU/m³ for houses according to CEC/Report #12 (Biological Particles in Indoor Environment) in the 1993 11, 47, 48. Rostami et al. (2017) conducted a study on the airborne fungi in indoor and outdoor air of the educational, treatment, research center, Ear, Nose and Throat ward, and hematology ward. Results showed that ENT ward for outdoor air (86 CFU) and hematology ward for indoor air (33 CFU) had the highest levels of airborne fungi ⁴⁹. Hoseinzadeh et Al. studied Dariai-e-Mehr DCC and found that the minimum and maximum fungal agents were 322 CFU/m³ and 1314 CFU/m³, respectively, which were higher than the results of this study ⁶.

Mentese et al. in Ankara found that the highest level of fungal contamination was observed in the kitchens of DCCs, preschools, restaurants, high schools, and homes, while Penicillium spp., Aspergillus spp., and Cladosporium spp. were the most observed fungi. A significant difference was found between the growth of fungi in the indoor and outdoor environment 50. The results of this study are consistent with our research. In other words, the highest frequency of fungi in our study was related to the fungal species of Aspergillus, Penicillium, and Cladosporium. Penicillium spores and Aspergillus are released in the environment more easily than Cladosporium and this is one of the reasons for the abundance of Penicillium species in the indoor air of children DCCs 51. Moreover, the distribution of Cladosporium particles is divided into larger particles than the Penicillium, which is more difficult to release the spores ⁵². Immonen et al. in a study in Finland on the indoor air of schools found that the most frequent species of fungi were Penicillium, Cladosporium, Aspergillus, and yeasts 51.

Aspergillus species, including Aspergillus flavus and Aspergillus niger were also observed in a study, where Aspergillus niger had the highest frequency and caused a variety of respiratory problems and allergic diseases 44. Chytridiomycota, such as Aspergillus, Penicillium, Cladosporium, and Alternaria (formerly known as Deuteromycota) are the largest group of fungi for reproduction, which cause allergic symptoms 53.

Aspergillus niger can cause angiomyceses, peritonitis, and endocarditis. It can also causes infection of the inner and middle ear as well as the

pulmonary aspergillosis ^{54, 55}. Another abundant species scattered in the environment is *Cladosporium*. In the presence of appropriate RH, these species can easily grow on the surfaces, which are allergic and may cause asthma in children with respiratory problems. In addition, the risk of spreading volatile organic compounds and mycotoxins is high by this species. Fungi can also produce secondary metabolites (fungal VOCs and mycotoxins) and the production of mycotoxins depends on the substrates where mould grows. This is also true for VOCs with lower degrees ^{56, 57}.

In public buildings, including DCCs, *penicillium* species are found easily in the air and particles and can grow under favorable surface moisture conditions ⁴³. In short, the indoor air is affected by the outdoor air; so, fungal species in the outdoor areas of the DCC can grow in the indoor environment if the conditions are favorable. For example, the *Penicillium* species typically exist in the outdoor air, but it grows easily under conditions in the indoor air ⁵⁸. Therefore, it is necessary to monitor the respiratory diseases, density and fungi concentration, as well as the personal health.

In the study carried out by Hwang in the city of Ulsan in South Korea on the seasonal concentration of airborne fungal load in children DCCs, the number of fungi was 0-1888 MPN/m³. The highest number of fungi was reported in the summer, while the lowest number was found in the winter. *Penicillium* and *Aspergillus* species were extracted from the cultured samples ⁵⁹. This study was consistent with the fungi found in our study, but it was different in the sampling seasons.

In a study conducted by Karwowska in elementary schools and high schools, the researchers found that the amount of contamination was significantly higher than EU standards; so that the amount of fungal contamination was estimated as 30-785 CFU/m³. The most frequent fungi were *Aspergillus* and *Penicillium* ⁶⁰. In a study by Hoseinzadeh et al., 30 CFU/m³ was considered as the standard to compare the results; the rates lower than 30 CFU/m³ were in accordance with the standard and values higher than 30 CFU/m³ were

1085

higher than the permissible limit ¹¹.

Sławomira et al. studied the concentration of bacteria and respiratory air fungi in five schools in Lublin, Poland and found that the concentration of a species of fungus was greater than the standard level (300 CFU/m³), equivalent to 690 CFU/m³ only in one of the schools. Among the fungal species, *Aspergillus* and *Penicilinium* had the highest arte ⁶¹. Meyer et al. in Finland studied molds in the indoor air of school buildings and found that some species of *Aspergillus* and *Penicillium* molds were the most important fungal species isolated from these sites ⁶². Their results are in the same line with those of our study.

Comparison of the selected samples from outdoor and indoor spaces indicated that outdoor samples were significantly more contaminated with Rhizopus fungus than indoor samples (P-value = 0.001). No relationship was found between contamination with other fungal agents and open or closed spaces, with the exception of Rhizopus. The results are presented in Table 3. Hosseinzadeh et al. investigated the indoor fungal bioaerosols of DCCs in Noor Abad City of Lorestan province in 2017. They showed that the most fungal species isolated from these environments was Penicillium type ⁶. The similarity between the results of these two studies can be attributed to the easier release of Penicillium and Aspergillus spores than Cladosporium spores. In this study, Pennicillium and Aspergillus flavus were the most common species in outdoor environment.

The results of this study showed that the I/O ratio was greater than 1 in all samples. So, presence of people and decreased rate of ventilation to save the energy or lack of proper ventilation can decrease the indoor air quality and increase the concentration of fungal bioaerosols in the environment ⁶³. This ratio indicates that the higher rate of indoor bioaerosols fungi, compared with outdoor environment, and is not consistent with the study of indoor/outdoor air quality for elementary schools in Lisbon ⁶⁴. In the study by Faridi et al., I/O rates were more than 1 and similar to the present study and the most frequent species were *Cladosporium*, *Penicillium*, and *Aspergillosis*

⁶³. In another study, the effects of the indoor and outdoor air were compared and the researcher concluded that the microbial samples and species obtained were very similar to those of outdoor environment ⁶⁵. In another study in DCCs, no difference was observed between the fungal contamination in the outdoor and indoor air ⁶⁶. However, we found that the contamination of the indoor environments was higher than the outdoor contamination.

Moreover, a study in the city of Lisbon on the quality of indoor and outdoor air of the elementary schools showed that the level of bacterial and fungal contamination was high and the factors influencing this amount were RH, T, wind speed, student density, and human activity ⁶⁴. The similarity and frequency of the airborne fungi in DCCs in public places can be attributed to the congestion of children and students in closed environments and the lack of proper ventilation ⁶.

Conclusion

This study was carried out in 10 DCCs of Khorramabad city. The results of 180 fungal samples showed that 96.1% of the plates had fungal colonies and 3.9% had no colony. The results showed *Fusarium* fungus with prevalence of 6% was only observed in spring (P-value = 0.014), while contamination with other fungal agents did not show a significant relationship with the target season.

The I/O ratio was more than 1, indicating that the indoor bioaerosols were higher than those of the outdoor environment.

Except for *Rhizopus*, no significant relationship was found between the concentration of fungal bioaerosols in the indoor and outdoor air. *Aspergillus niger* and *Mucor* were the most abundant species in the indoor air and the outdoor fungal agents were similar to the indoor air samples. Furthermore, *penicillium* species with 31.1 % and *Aspergillus flavus* with 21.7% were the most frequent fungi. A significant correlation was found between the ambient temperature, RH, and the observed fungal colony count.

The results indicated that, the central and

standard ventilation system in the DCCs was not used for air conditioning and natural air was used for ventilation without treatment. In addition, the high level of bioaerosol in the air of DCCs can be due to the large number of children, lack of using personal and non-sterile items by children, lack of having a suitable ventilation system, and improper building design of the DCCs. Moreover, the contamination of these sites can be due to the lack of suitable health standards in the field of educator traffic and parents, the ratio of children number to the capacity of the class, and the lack of guidelines or standards in Iran to monitor the fungal agents in DCCs. Therefore, some recommendations are suggested as follows: 1. Using a ventilated system equipped with a filtration system (the DCCs did not have air treatment systems), 2. Modifying the ventilation system and buildings of DCCs, 3. Monitoring the fungal contaminants continuously, and 4. Reducing the population of children in closed environment and controlling the standard number of children in the class room.

Limitations of the study

In the sampling sites, a single point was selected and monitored. However, using more sampling points could cover the actual bioaerosol concentration much better in the results. In this study only culturable and viable fungi bioaerosols were assessed. We suggest the future researchers to study non-viable and non-culturable bioaerosls such as viruses, pollens, cellular fragments, and toxins in future similar studies. In the present study characterization of bioaerosols was performed up to the genus level. In order to identify the bioaerosols, more detailed flow cytometry and molecular techniques are recommended.

Acknowledgment

The authors would like to thank the DCCs of Khorramabad for their help and Nasrin Shokrpour from Shiraz University of Medical Sciences for English editing of this article.

Funding

This study was supported by Islamic Azad University of Ahvaz (Grant number of thesis is 10650516942009).

Conflict of Interest

The authors of this study declare that they have no conflict of interests.

This is an Open-Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt, and build upon this work, for commercial use.

References

- 1.Mandal J, Brandl H. Bioaerosols in indoor environment-a review with special reference to residential and occupational locations. The Open Environmental & Biological Monitoring Journal. 2011;4(1):83-96.
- 2.Georgakopoulos D, Després V, Fröhlich-Nowoisky J, et al. Microbiology and atmospheric processes: biological, physical and chemical characterization of aerosol particles. Biogeosciences. 2009;6(4):721-37.
- 3.Humbal C, Gautam S, Trivedi U. A review on recent progress in observations, and health effects of bioaerosols. Environ Int. 2018; 118:189-93.
- 4.Dedesko S, Stephens B, Gilbert JA, et al. Methods to assess human occupancy and occupant activity in hospital patient rooms. Build Environ. 2015;90:136-45.
- 5.Cui W, Cao G, Park JH, et al. Influence of indoor air temperature on human thermal comfort, motivation and performance. Build Environ. 2013;68:114-22.
- 6.Hoseinzadeh E, Taha P, Sepahvand A, et al. Indoor air fungus bioaerosols and comfort index in day care child centers. Toxin Rev. 2017; 36(2):125-31.
- 7.Lee AK, Lau AP, Cheng JY, et al. Source identification analysis for the airborne bacteria and fungi using a biomarker approach. Atmos Environ. 2007;41(13):2831-43.
- 8.Lang-Yona N, Dannemiller K, Yamamoto N, et al. Annual distribution of allergenic fungal spores in atmospheric particulate matter in the Eastern Mediterranean; a comparative study between ergosterol and quantitative PCR analysis. Atmos Chem Phys. 2012;12(5):2681-

100/

90.

- 9. Aydogdu H, Asan A. Airborne fungi in child day care centers in Edirne City, Turkey. Environ Monit Assess. 2008;147 (1-3): 423-44.
- 10. Stępalska D, Wołek J. Variation in fungal spore concentrations of selected taxa associated. Aerobiologia (Bologna). 2005;21(1):43-52.
- 11. Hoseinzadeh E, Samarghandie MR, Ghiasian SA, et al. Evaluation of bioaerosols in five educational hospitals wards air in Hamedan, During 2011-2012. Jundishapur J Microbiol. 2013;6(6):e10704.
- 12. Branco P, Alvim-Ferraz M, Martins F, et al. The microenvironmental modelling approach to assess children's exposure to air pollution—a review. Environ Res. 2014;135: 317-32.
- 13. Branco P, Alvim-Ferraz M, Martins F, et al. Children's exposure to indoor air in urban nurseries-part I: CO2 and comfort assessment. Environ Res. 2015;140:1-9.
- 14. Kabir E, Kim K-H, Sohn JR, et al. Indoor air quality assessment in child care and medical facilities in Korea. Environ Monit Assess. 2012;184(10):6395-409.
- 15. Qu F, Weschler LB, Sundell J, et al. Increasing prevalence of asthma and allergy in Beijing pre-school children: Is exclusive breastfeeding for more than 6 months protective? Chin Sci Bull. 2013;58(34):4190-202.
- 16. Vijayalakshmi I, Rao PS, Chugh R. A comprehensive approach to congenital heart diseases: JP Medical Ltd; 2013.
- 17. Madureira J, Paciência I, Rufo JC, et al. Assessment and determinants of airborne bacterial and fungal concentrations in different indoor environments: Homes, child day-care centres, primary schools and elderly care centres. Atmos Environ. 2015;109:139-46.
- 18. Ciucci E, Calussi P, Menesini E, et al. Seasonal variation, weather and behavior in daycare children: a multilevel approach. Int J Biometeorol. 2013;57(6):845-56.
- 19. Roda C, Barral S, Ravelomanantsoa H, et al. Assessment of indoor environment in Paris child day care centers. Environ Res. 2011;111(8):

1010-7.

- 20. Fernández LC, Alvarez RF, González-Barcala FJ, et al. Indoor air contaminants and their impact on respiratory pathologies. Arch Bronconeumol. 2013;49(1):22-7.
- 21. Rao D, Phipatanakul W. Impact of environmental controls on childhood asthma. Curr Allergy Asthma Rep. 2011;11(5):414-20.
- 22. Fard R, Hosseini M, Faraji M, et al. Building characteristics and sick building syndrome among primary school students. Sri Lanka Journal of Child Health. 2018;47(4):332-7.
- 23. Heseltine E, Rosen J. WHO guidelines for indoor air quality: dampness and mould: WHO Regional Office Europe; 2009.
- 24. Ghaderpoori M. Heavy metals analysis and quality assessment in drinking water—Khorramabad city, Iran. Data Brief. 2018;16: 685.
- 25. Nourmoradi H, Goudarzi G, Daryanoosh SM, et al. Health impacts of particulate matter in air using AirQ model in Khorramabad city, Iran. Journal of basic research in medical sciences. 2015;2(2):44-52.
- 26. Goh I, Obbard J, Viswanathan S, et al. Airborne bacteria and fungal spores in the indoor environment. A case study in Singapore. Acta Biotechnol. 2000;20(1):67-73.
- 27. Gołofit-Szymczak M, Górny RL. Bacterial and fungal aerosols in air-conditioned office buildings in Warsaw, Poland-he winter season. Int J Occup Saf Ergon. 2010;16(4):465-76.
- 28. Naddafi K, Hassanvand MS, Yunesian M, et al. Health impact assessment of air pollution in megacity of Tehran, Iran. Iranian J Environ Health Sci Eng. 2012;9(1):28.
- 29. Hameed AA, Khoder M, Yuosra S, et al. Diurnal distribution of airborne bacteria and fungi in the atmosphere of Helwan area, Egypt. Sci Total Environ. 2009;407(24):6217-22.
- 30. Hsu Y-C, Kung P-Y, Wu T-N, et al. Characterization of indoor-air bioaerosols in Southern Taiwan. Aerosol Air Qual Res. 2012;12:651-61.
- 31. Hoseini M, Jabbari H, Naddafi K, et al. Concentration and distribution characteristics of

- airborne fungi in indoor and outdoor air of Tehran subway stations. Aerobiologia (Bologna). 2013;29(3):355-63.
- 32. Wang Y-F, Wang C-H, Hsu K-L. Size and seasonal distributions of airborne bioaerosols in commuting trains. Atmos Environ. 2010;44(35): 4331-8.
- 33. Scaltriti S, Cencetti S, Rovesti S, et al. Risk factors for particulate and microbial contamination of air in operating theatres. J Hosp Infect. 2007;66(4):320-6.
- 34. Aneja K. Experiments in microbiology, plant pathology and biotechnology. New Age International: 2007.
- 35. Fang Z, Ouyang Z, Hu L, et al. Culturable airborne fungi in outdoor environments in Beijing, China. Sci Total Environ. 2005;350(1-3):47-58.
- 36. Wang W, Ma Y, Ma X, et al. Seasonal variations of airborne bacteria in the Mogao Grottoes, Dunhuang, China. Int Biodeterior Biodegradation. 2010;64(4):309-15.
- 37. De Hoog G, Guarro J, Gené J, et al. Atlas of clinical fungi. 2000. Utrecht/Reus, The Nederlands: 201.
- 38. Ellis DH, Davis S, Alexiou H, et al. Descriptions of medical fungi: University of Adelaide Adelaide; 2007;61:167.
- 39. Fisher F, Cook N. Fundamentals of diagnostic mycology. 1998. WB Saunders Philadelphia, PA.
- 40. Handbook A. HVAC applications. ASHRAE Handbook, Fundamentals. 2007.
- 41. Frankel M, Bekö G, Timm M, Gustavsen S, Hansen EW, Madsen AM. Seasonal variations of indoor microbial exposures and their relation to temperature, relative humidity, and air exchange rate. Appl Environ Microbiol. 2012;78(23):8289-97.
- 42. Moungthong G, Klamkam P, Mahakit P, et al. Efficacy of the Precise Climate Controller on the reduction of indoor microorganisms. Asia Pac Allergy. 2014;4(2):113-8.
- 43. Tang JW. The effect of environmental parameters on the survival of airborne infectious agents. J R Soc Interface. 2009;6(Suppl 6):S737-S46.

- 44. Aquino S, de Lima JEA, do Nascimento APB, et al. Analysis of fungal contamination in vehicle air filters and their impact as a bioaccumulator on indoor air quality. Air Qual Atmos Health. 2018;11(10):1143-53.
- 45. Park D, Jo K, Yoon C, et al. Factors influencing airborne concentration of fungi, bacteria and gram negative bacteria in kindergarten classroom. Korean Journal of Environmental Health Sciences. 2004;30(5):440-8.
- 46. Sepahvand A, Shams-Ghahfarokhi M, Allameh A, et al. A survey on distribution and toxigenicity of Aspergillus flavus from indoor and outdoor hospital environments. Folia Microbiol (Praha). 2011;56(6):527-34.
- 47. Castleman BI, Ziem GE. American conference of governmental industrial hygienists: Low threshold of credibility. Am J Ind Med. 1994;26(1):133-43.
- 48. Rao CY, Burge HA, Chang JC. Review of quantitative standards and guidelines for fungi in indoor air. J Air Waste Manag Assoc. 1996;46(9):899-908.
- 49. Rostami N, Alidadi H, Zarrinfar H, et al. Assessment of indoor and outdoor airborne fungi in an Educational, Research and Treatment Center. Italian journal of medicine. 2017;11(1):52-6.
- 50. Mentese S, Arisoy M, Rad AY, et al. Bacteria and fungi levels in various indoor and outdoor environments in Ankara, Turkey. Clean (Weinh). 2009;37(6):487-93.

- 51. Immonen J, Meklin T, Taskinen T, et al. Skinprick test findings in students from moisture-and mould-damaged schools: A 3-year follow-up study. Pediatr Allergy Immunol. 2001;12(2):87-94.
- 52. Hoseinzadeh E, Samarghandi MR, Ghorbani Shahna F, et al. Isoconcentration mapping of particulate matter in H amedan intercity bus stations. Water Environ J. 2013;27(3):418-24.
- 53. Cerqueira P, Guimarães A. Indoor air quality in a petrochemical industry. Cientefico. 2017; 17(35):1-18.
- 54. Gupta VK. New and future developments in

- microbial biotechnology and bioengineering: Aspergillus system properties and applications: Elsevier; 2016 Oct 27.
- 55. Moazam S, Denning DW. Aspergillus nodules in chronic granulomatous disease attributable to Aspergillus ochraceus. Med Mycol Case Rep. 2017;17:31-3.
- 56. Vacher G, Niculita-Hirzel H, Roger T. Immune responses to airborne fungi and non-invasive airway diseases. Semin Immunopathol. 2015;37(2):83-96.
- 57. Yang CS, Johanning E. Airborne fungi and mycotoxins. Manual of Environmental Microbiology, 3rd ed. American Society of Microbiology; 2007.
- 58. Sinclair R, Gerba CP. Microbial contamination in kitchens and bathrooms of rural Cambodian village households. Lett Appl Microbiol. 2011;52(2):144-9.
- 59. Hwang K, Lee A-M, Shin H, et al. Seasonal monitoring of airborne microbial concentrations in kindergatens. Korean Journal of Microbiology. 2003;39(4):253-9.
- 60. Karwowska E. Microbiological air contamination in some educational settings. Pol J Environ Stud. 2003;12(2):181-6.

- 61. Dumała SM, Dudzińska MR. Microbiological indoor air quality in Polish schools. Annual Set The Environment Protection (Rocznik Ochrona Środowiska). 2013;15:231-44.
- 62. Meyer HW, Würtz H, Suadicani P, et al. Molds in floor dust and building-related symptoms in adolescent school children. Indoor Air. 2004;14(1):65-72.
- 63. Faridi S, Hassanvand MS, Naddafi K, et al. Indoor/outdoor relationships of bioaerosol concentrations in a retirement home and a school dormitory. Environ Sci Pollut. 2015; 22(11):8190-200.
- 64. Pegas P, Alves C, Evtyugina M, et al. Indoor air quality in elementary schools of Lisbon in spring. Environ Geochem Health. 2011;33(5): 455-68.
- 65. Adams RI, Bhangar S, Pasut W, et al. Chamber bioaerosol study: outdoor air and human occupants as sources of indoor airborne microbes. PLoS One. 2015;10(5):e0128022.
- 66. Shin S-K, Kim J, Ha S-m, et al. Metagenomic insights into the bioaerosols in the indoor and outdoor environments of childcare facilities. PLoS One. 2015;10(5):e0126960.