

Journal of Environmental Health and Sustainable Development



Microbial Community Analysis Using MiSeq Sequencing and Pathway of Methane Production in Tehran WWTP: A Full-Scale Anaerobic Digester

Reza Barati Rashvanlou^{1,2}, Mahdi Farzadkia^{1,2}, Abbas Ali Moserzadeh³, Asghar Riazati⁴, Chiang Wei⁵, Edris Hoseinzadeh⁶*

- ¹ Research Center for Environmental Health Technology, Iran University of Medical Sciences, Tehran, Iran.
- ² Department of Environmental Health Engineering, School of Public Health, Iran University of Medical Sciences, Tehran, Iran.
- ³ Department of Civil Engineering, Amirkabir University of Medical Sciences, Tehran, Iran.
- ⁴ Managing Director of Tehran Water and Wastewater Co., Tehran, Iran.
- ⁵ Research Fellow, the Experimental Forest, College of Bio-Resources and Agriculture, National Taiwan University, Nantou 55750, Taiwan.
- ⁶ Student Research Committee, Saveh University of Medical Sciences, Saveh, Iran.

ARTICLE INFO

ORIGINAL ARTICLE

Article History:

Received: 05 May 2020 Accepted: 10 July 2020

*Corresponding Author:

Edris Hoseinzadeh

Email:

edris-2009@live.com

Tel:

+988642343395

Keywords:

Microbial Community, Methane Production Pathway, Biogas, Anaerobic Digester.

ABSTRACT

Introduction: One of biological wastewater treatment methods that utilizes to both digesting waste activated sludge and methane production is anaerobic digestion (AD). It is believed to be most effective solution in terms of energy crisis and environmental pollution issues.

Materials and Methods: In this study the sludge was digested anaerobically sampled from a full-scale WWTP, located at south of Tehran, Iran for evaluation. To study the microbial community within the sludge the MiSeq Sequencing method utilized. Based on our field data and microbial community data, a schematic diagram of probable leading pathways was made in the studied digester.

Results: At first, the community variety in the bulk sludge and richness were enhanced followed by loading increasing. Meanwhile, the loading change enhanced the community richness and variety of the sludge. By comparing the rank-abundance distributions, a shallow gradient would show high evenness since the abundances of diverse species are alike. The results showed all the communities were extremely diverse and 15 phyla were distinguished in the sludge sample. The dominant phyla of the community were *Bacteroidetes* and *Firmicutes* and quantity of the two phyla were 21% and 11%, respectively. *Anaerobaculum, Acinetobacter, Syntrophomonas,* and *Coprothermobacter* were the chief genera for the microbial communities and the sum of four genera were 7%, 3%, 3%, and 2%, respectively.

Conclusion: It was shown that syntrophic acetate oxidizing bacterias (SAOBs) metabolized acetate through hydrogen trophic methanogenesis in the digester. Generally, the findings may be useful to help the wastewater operators to utilize an effective method that able to treat waste sludge plus methane production, simultaneously.

Citation: Barati Rashvanlou R, Farzadkia M, Moserzadeh AA, et al. *Microbial Community Analysis Using MiSeq Sequencing and Pathway of Methane Production in Tehran WWTP: A Full-Scale Anaerobic Digester.* J Environ Health Sustain Dev. 2020; 5(3): 1091-102.

[DOR: 20.1001.1.24766267.2020.5.3.8.7]

Introduction

Activated sludge process uses a biological process to treat municipal and industrial wastewater ¹. This process is noticed by process designers due to its high efficiency in removing/decreasing most pollutants established as the discharge standards to acceptable levels ^{2, 3}. The wastewater treatment plant operation reports that the sludge generation through biological wastewater treatment is the main activated sludge side effects because this process needs more than 50-60% of the integrated wastewater management budget 4. The activated sludge process should be moderated to decrease sludge generation. Anaerobic digestion is highly efficient in sludge treating and bioenergy generating simultaneously 5-7. By applying this process, the treated sludge had acceptable organic matter, pathogens, and nutrient content for use in land application. In addition, the sludge quantity decreased and led to easy final management and generated methane as a valuable biogas. Generally, the anaerobic digestion process is capable of converting 50% bio-solids (in average) to methane 8. The microbial communities in the sludge are the heart of efficient methane generation through anaerobic digestion and sludge reduction. In other words, we need to have enough information about the sludge microbial community to have a selective pathway to operate and generate methane. Furthermore, if some is predominant, microbial community pathways change that leads to unpleased results. Generally, assessing the microbial community of sludge is one of the necessary steps to have an efficient and cheap anaerobic sludge digestion process 9, 10. Based on the available reports, some molecular approaches exist to find predominant microbial communities in anaerobic systems. Among them, Illumina sequencing and 454 pyrosequencing technologies at high-throughout sequencing method category are the most efficient user-friendly methods microbial and community analysis anaerobic digested sludge 11-¹³. By exploring the literature, we found the Illumina sequencing as the most interesting method to assess microbial communities among soil, ocean, human gut microbes, and activated sludge 14, 15. The use of Illumina sequencing to assess anaerobic sludge community is rare, but its application is rising. By in-depth analysis of the microbial community structure, the microbial functions can be identified in anaerobic sludge digestion and methane production pathway. The present study aimed to explore the microbial community arrangement in a full-scale anaerobic digester of Tehran wastewater treatment plant. To this end, DNA extraction was done from a fullscale anaerobic digestion sludge. The dominant microorganisms, functional profiles, methane production pathway from AD of sludge investigated. were Especially, significant microorganisms involved in sludge digestion methane production were analyzed and comprehensively according to the acquired molecular data. This study aimed to assess the microbial community structure of a full-sized wastewater treatment having AD for sludge managing and methane production, simultaneously.

Materials and Methods

Sampling of AD sludge from a full-scale **WWTP**

In this study a full-scale WWTP, located at south of Tehran, Iran selected. There is used AD to manage activated sludge. The south wastewater treatment plant of Tehran is located in a region of about 110 hectares in the south of Rey. The plan is envisioned for the population of 2,100,000 people. The total capacity of the plant is 5.2 m³/s having four modules. Wastewater treatment method is activated sludge conventional system. The extra sludge from the primary sedimentation goes into gravity sludge thickening. Later, the excess sludge from the secondary sedimentation was thickened using a mechanical thicker and went into the sludge digester. The studied fullscale anaerobic reactor digesting activated sludge including thickening tanks, anaerobic mesophilic digestion and dewatering. Figure 1 presents wastewater treatment process diagram.




Figure 1: representation of the elements of the studied full-scale WWTP and the sampling location (as demonstrated red stars); the wastewater plant section is as following: 1) main sewer; 2) mail sewer; 3) screen; 4) grit and grease removal; 5) primary sedimentation; 6) high loaded biological stage (C- removal and partial N-removal); 7) aeration blower; 8) secondary sedimentation; 9) return sludge; 10) tricking filter; 11) disinfection; 12) main outlet; 13) primary sludge thickener; 14) belt thickener; 15) sludge digester; 16) desulphurization; 17) excess gas burner; 18) gas storage; 19) generator; 20) sludge thickening; 21) sludge dewatering.

DNA extraction, library construction and sequencing

At first, the sampled sludge mixed with ethanol (with purity of ~100%) at 1:1 ratio (v/v %). Afterward, it was transferred to lab in standard condition (using a cool box with -20 °C temperature). All the qualified DNAs (that were checked by quality test) were applied to construct libraries. Gel electrophoresis (1% agarose gel, Voltage: 150 V; Electrophoresis Time: 40 min) was applied to evaluate the quality of

used to specify the concentrations of DNA. The DNA concentration of the sample was 52.6 ng/ μ L. The microbial community sequencing was done with Illumina Hi Seq/Mi Seq platform (by NIGEB Genome Center, China). Meta rDNA Amplicon sequencing includes Meta rDNA V3, V6, V4, V1-V3, V3-V4, ITS1, and ITS2 regions amplicon library construction. The paired-end reads with overlap were merged to tags and the tags were clustered to the Operational Taxonomic Unit (OUT) at 97% sequence similarity by scripts of USEARCH software (v7.0.1090). The library was

sequenced on an Illumina Hi Seq/Mi Seq platform. If the two paired-end reads overlapped, the consensus sequence would be generated by FLASH (Fast Length Adjustment of Short reads, v1.2.11). The parameters adopted for overlapping were the minimal overlaping length of 15 bp and the allowed mismatching ratio of overlapped region was 0.1.

Jehsd.ssu.ac.ir

1093

Microbial Community analysis

USEARCH software (v. 7.0.1090) scripts were used to cluster the tags into OTU. The tags were clustered into OTU with a 97% threshold using UPARSE and the OTU unique representative sequences were obtained. UCHIME (v4.2.40) was used to filter out the chimeras. By matching the gold database (V20110519, UNITE (V20140703)) with 16S rDNA and ITS sequences distinctly, de novo chimera detection was done for 18S rDNA sequences. All tags were mapped to each OTU representative sequences using **USEARCH** GLOBAL and then the number of tags for each OTU in each sample was summarized to OTU

abundance table. Ribosomal Database Project (RDP) Classifier (v. 2.2) trained on the Greengenes database was used along with 0.8 confidence values as cutoff to classify OTU representative sequences taxonomically. The aligned DNA sequences were searched using the Silva database (Silva (default): V119; http://www.arb-silva.de) etc. 18S rDNA was used for fungal community to identify each OTU. The ITS was also used for fungal community (UNITE (default): Version. 6 20140910). In addition, 16S rDNA was used for bacterial and Archaea community (Greengene (default): V201305; RDP: Release 9 201203). Taxonomy cranks were assigned to OTU representative sequence using Ribosomal Database Project (RDP; http://www.rdp.cme.msu.edu/) Na e Bayesian Classifier v.2.2. Finally, OTU and taxonomic ranks were used to analyze the alpha diversity, beta diversity, and different species screening.

Species phylogenetic and Diversity analysis

The Silva core set (Silva_108_core_ aligned_seqs) using PyNAST by 'align_seqs.py' used to represent lined up sequences. The QIIME (v1.80) built-in scripts were used to construct the representative OTU phylogenetic tree. The tags with a maximum abundance of each Genus were selected as the matching Genus-representative sequences. In addition, the Genus level phylogenetic tree was obtained the same as OTU phylogenetic tree. R (v3.0.3) software was used to draw the phylogeny

tree. The rarefaction curve was drown using the indices calculated with extracted tags of the samples. Mothur (v1.31.2) was applied to calculate the indices and the corresponding rarefaction curve was drawn by R (v3.0.3) rarefaction curve. Each index was formulated and calculated by the following address: http://www. mothur.org/wiki/ Calculators. Furthermore, the microbial diversity indices were Chao-the Chao1 estimator, ACE-the ACE estimator, Shannon- the Shannon index, Simpson- the Simpson index, and goods coverage index that can be calculated using the following addressws, respectively (http://www.mothur. org/ wiki/Chao). (http://www. mothur.org/wiki/Ace), (http:// www. mothur.org/ wiki/Shannon), and (http://www. mothur. org/ wiki/ Simpson, (http:// www. mothur. org/ wiki/ Coverage). To analyze the complexity of the species' diversity, the Alpha diversity was used (http://www.mothur.org/wiki/ Schloss_SOP#Alpha_diversity).

Results

Microbial Richness and Diversity Evolution

Microbial community analysis performed by MiSeq sequencing method. The diversity estimators of Chao1, ACE, and Shannon index, Simpson index, and P-value of the sequencing for each sample are shown in Table 1.

Rarefaction, Rank-abundance distributions

As shown in figure 2, the rarefaction curves were generated for the sludge sample.

Table 1: The resluts for species richness, phylogenetic diversity and the microbial community in the studied sludge (ACE richness estimator; Chaorichness estimator; Shannon diversity index; Simpson diversity index).

Sample	Sobs	R	ichness	Diversity		
Sample		Chao	Ace	Shanon	Simpson	
1	491	491	491	3.597944	0.070040	
2	191 ± 43.24	292.44 ± 16.15	382.62 ± 177.89	2.33 ± 4.34	0.18 ± 0.036	
3	682.5 ± 166.91	898.39 ± 154.11	854.80 ± 154.54	3.04 ± 4.27	0.15 ± 0.04	
P value	0.00075	0.00065	0.00114	0.00066	0.00167	

[DOI: 10.18502/jehsd.v5i3.4280

Jehsd.ssu.ac.ir




Figure 2: Different curves based on the observed species value, chao1 value, ACE value, shannon value, and simpson value

Microbial Phylum and Class Identification

In order to find the phyla and class of microbial community the confirmed reads used. represented in figure 3, all communities were very diverse and 15 phyla were noticed in the sludge sample. The key phyla that was found in the studied sludge sample were Bacteroidetes and Firmicutes, while the sums of phyla abundances were 21% and 11% in the respective order. Furthermore, sequences belonging $(\leq 1\%)$ Actinobacteria $(\leq 1\%),$ Acidobacteria *Armatimonadetes* (6%), BRC1(≤1%), *Chlamydiae* $(\leq 1\%)$, Chloroflexi $(\leq 2\%)$, Fusobacteria $(\leq 1\%)$, Planctomycetes ($\leq 1\%$), Proteobacteria (7%), Spirochaetes $(\leq 1\%),$ Synergistetes (7%),Thermotogae (6%), and Verrucomicrobia $\leq 1\%$ were identified. The representative sequences were aligned against the Silva core (Silva_108_core_aligned_seqs) using PyNAST by 'align_seqs.py'. A representative **OTU** phylogenetic tree was constructed using the QIIME (v1.80) built-in scripts including the fast tree method for tree construction. The tags with the highest abundance of each Genus were chosen as the corresponding Genus representative sequences and the Genus level phylogenetic tree was obtained similar to OTU phylogenetic tree. The phylogeny tree was imaged by R (v3.0.3) software at last.

Genus species phylogeny tree




Figure 3: Genus level phylogenetic tree (The same Phylum is shown as the same color)

Microbial Genera Evolution

According to our findings (Figure 4), there were different microbial genera in the studied sludge. The bacteria belonging to the genera *Anaerobaculum*, *Acinetobacter*,

Syntrophomonas, Coprothermobacter, Clostridium,

Gp 10, Lutispora, Proteiniborus, Sedimentibacter, and Soehngenia were prevailing in the sludge. Anaerobaculum, Acinetobacter, Syntrophomonas, and Coprothermobacter were the key genera for the microbial communities and the sum of four genera were 7%, 3%, 3%, and 2%, respectively.




Figure 4: The taxonomics composition distribution histograms of samples at Phylum, Order, Class, Family, Genus, and Species level separately. At Phylum, all species were used to draw the histogram. The species with an abundance of less than 0.5% in all samples were classified into 'others'.

Discussion

In this study, the microbial community within a full-Scale AD sludge determined using the MiSeq sequencing method. Meanwhile, the pathways of methane production in the studied WWTP were proposed.

Microbial Richness and Diversity Evolution

Logged species value, chao1 value, and ACE value can reflect the species' richness of the community. It is possible to use the rarefaction curve according to the three values to assess if produced data is sufficient for covering all species in the community 16, 17. The community richness of the sludge had diverse values based on the community diversity of sludge. Rationally, the increased loading or changing the sludge property that may be as a result of alternation in wastewater biodegradable organic matter content, can affect the diversity and richness in the sludge, considerably 18. The communities of sludge showed a minor decrease in richness and diversity for sample two. According to table 1, the highest richness and diversity was logged for sample three. According to our results, the species richness and phylogenetic diversity increased for the bulk sludge; then, it reduced by increase of loading.

Rarefaction, Rank-abundance distributions

The OTU rank abundance curve makes it possible to show the species' richness and species' evenness visually¹⁷. According to figure 4, the species are ranked based on their richness. By obtaining the slope of fitted line, the species' evenness was counted. A steep gradient is an indication of low evenness¹⁹. The comparison of rank-abundance distributions reveals a shallow gradient highlighting high evenness since the abundances of various species are similar. When microbiota are gathered from a certain niche, it is encessary to assess how well a sample shows the true diversity of the specific niche. In other wods, it is identical with species' richness and relative abundance in time and space. Rarefaction curves measure the OTUs observed with a given depth of sequencing applied for comparing the observed richness among communities with an unequal samples¹⁷. As the table 1 shows the rarefaction result is in-line with Chao1, ACE, Shannon, and Simpson results. It can be concluded that the species richness and phylogenetic diversity have a similar alternation arrangement.

Regarding higher microbial diversity in the sludge cause enhancing microbial activity, so organic loading rate increasing unable to have considerable effect on Chemical Oxygen Demands (COD) efficiency ^{16, 20}. Alternation of sludge microbial community (Quantitatively and

1098

that can be (HRT) and Here, HRT similar; how flactuation be Microbial For bacter recurrently many anaer

qualitatively) can be result of environmental steress that can be caused by hydraulic retention time (HRT) and operation temperatures changes ^{21, 22}. Here, HRTs and operation temperatures were similar; however, the organic loading rate had flactuation because of wastewater property.

Microbial Phylum and Class Identification

For bacteria, Firmicutes and Bacteroidetes were recurrently recognized as the chief phyla in many anaerobic digesters²³. It is believed that Firmicutes and Bacteroidetes are significant in a hydrolytic/acidogenic digester fed with dried hay and straw²⁴. It was also reported that these two phyla are vital during batch digestion. The phylum Bacteroidetes is a group of bacteria found in many different habitats, which live in soil or aquatic environments. It is also related to humans and animals that possess a high level of phenotypic and metabolic diversity 25. Bacteroidetes are able to release more proteinaceous EPS that facilitate its colonization ^{16, 26}. Their fimbriae helps them adhesion to other bacteria cells (colony forming) and to solid surfaces and improving their enrichment on the surfaces.

Several other studies also offered Bacteroidetes as one of the key microbial components in anaerobic reactors²⁷. Bacteroidetes are fermentative bacteria that play a role in degradation of carbohydrates and proteins into acetate and NH₃. The second chief bacterial phylum - Firmicutes represented 11% of the total bacterial clones in sludge sample. Firmicutes are hydrolytic bacteria that are capable of producing extracellular enzymes such as cellulases, lipases, and proteases^{28, 29}. Firmicutes is involved in the hydrolysis step of the anaerobic digestion (AD) sludge process by metabolizing a wide variety of compounds into acetate and butyrate (important **VFA** intermediates)¹⁶. According to Ding et al., Bacteroidetes, Chloroflexi, and Candidate with majority of 73.29% were the dominant phyla of the M-AnMBR sludge. As shown in figure 4, the abundance of Bacteroidetes in the current study is similar to the one studied by Ding et al. However, Chloroflexi in our study was lower (2%). As

illustrated in figure 4, nine bacterial classes were detected in the sludge sample. Acidobacteria Gp 10, Anaerolineae, Bacteroidia, Betaproteobacteria, Clostridia, Deltaproteobacteria, Gammaproteobacteria, Synergistia, Thermotogae were the main classes. The Clostridia, Thermotogae, Synergistia, Gammaproteobacteria were the key four classes that accounted for 14%, 7.5%, 7%, and 5% of the total population in the sludge. Clostridia classes are bacilliform bacteria that respirate anaerobically. These class of bacterium was reported as one of the dominant bacterial class in the anaerobic sludge digesters¹⁶. The anaerobic sludge digestion can make a suitable environment for their growth ³⁰. The includes Thermotogae anaerobic. phylum thermophilic, as well as mesophilic bacteria. Some Thermotogae species are capable of surviving at high temperature. The bacteria of this class have metabolic ability to produce H₂ gas from complexcarbohydrates that makes them a likely candidate as a biotechnological source for CH₄ production³¹. Synergistia (phylum Synergistetes) that is one of the most abundant class of bacteria that are capable of decomposing complex organic matters efficiently. Spirochaetes is chemohetrotroph microorganism that have spiral shape. Regarding Spirochaetes can be found in AD reactor, so they able to recover methane through fermentation of carbohydrates or amino acids. Consequently, contribution of the above-mentioned three classes in the present study was possibly related to the organic matter removal within the digester. We also observed the Gammaproteobacteria is a class of the phylum Proteobacteria³². Some Gammaproteobacteria oxidize methane, photosynthetic, and hydrogen sulfide instead of water, producing sulfur as a waste product. Comparing our results with other systems with different in HRT and treating pattern showed that their microbial communities in the sludge were different with present study.

Microbial Genera Evolution

The literature reported high proportions of unidentified sequences at the genus level, which is confirmed in this study. Obviously, isolation and

cultivation of uncultured microorganisms in the AD process³³⁻³⁶ is a useful way to find out their role in the process, as recommended earlier. The genus Anaerobaculum, as the most overriding genus in genus level, comprises of Anaerobaculum thermoterrenum and Anaerobaculum mobile species and is a member of the phylum Synergistetes. The presence of Anaerobaculum strains in petroleum reservoirs, mining wastewater, solid waste digester, and methanogenic reactors were also reported. Anaerobaculum strains are fermentative organisms capable of producing acetate, CO2, and H2 as their major metabolic products³⁷. They are unable to reduce sulfate or sulfite to hydrogen sulfide. Sugihara et al. showed that Anaerobaculum-related species were dominant bacteria³⁸. They concluded that Anaerobaculum-related species were able to degrade propionate. Consequently, one can say that higher abundance of Anaerobaculum in the studied digester possibly led to no accumulation of propionic acid and H₂. According to the hydrolysis, acidogenesis, acetogenesis, and methanogenesis are the possible biogas production process stages³⁹. At three first stages, the bacteria are principal organisms. CH₄ is formed through decarboxylation of acetate and methanisation of CO2 and H2 by acetogenotrophic and hydrogenotrophic archaea. Besides, synergistic acetate oxidation is involved in biogas formation. Moreover, during the hydrolysis of oils and fats, long-chain fatty acids (LCFA) are produced that are normally found in fattywastewaters. It is possible to convert these

compounds effectively to methane in anaerobic

bioreactors by applying the suitable condition. The LCFA converts to acetate and hydrogen by synchronizing the activity of syntrophic bacteria. In addition, the methanogenic archaea that utilizes these substrates is activated in this pathway. By prevailing this pathway, the complete LCFA degradation can be obtained. Here, the genus identification showed that Syntrophomonas sapovorans is the second most abundant species in the sludge sample. Meanwhile, studies show that 7 species of syntrophic bacteria are capable of growing on LCFA with more than 12 carbon atoms. Among these bacteria, only three mesophilic species utilizing the capable unsaturated LCFA, namely Syntrophomonas sapovorans (there was in studied sludge digester), Syntrophomonas curvata, and Syntrophomonas zehnderi. Generally, Syntrophomonas sapovorans anaerobic, syntrophic and fatty acid-oxidizing, and obligately proton-reducing.

Based on our field data and microbial community data, a schematic diagram of the probable leading pathways can be made in the studied digester (Figure 5). Acetate was metabolized via hydrogenotrophic methanogenesis by syntrophic acetate-oxidizing bacteria (SAOBs) in the studied digester. This is supported by the high abundance of Anaerobaculum in the digesters⁴⁰, which was considered as syntrophic acetate oxidizers. Syntrophic acetate oxidation occurs under elevated temperatures and high ammonia concentrations, which was also represented in the studied digester (Table 2).

Table 2: The sludge digester supernatant analysis; TN: Total Nitrogen, TP: Total Phosphates, COD: Chemical Oxygen Demand, BOD₅: Biochemical Oxygen Demand, Alk: Alkalinity.

NH ₄ -N(mg/L)	TN (mg/L)	TP (mg/L)	$P - PO_4^{-3}$ (mg/L)	COD (mg/L)	BOD ₅ (mg/L)	Alk (mg/L)	pН
960	1520	47.7	46.5	2104	410	10580	7.47

Jehsd.ssu.ac.ir

1100

DOR: 20.1001.1.24766267.2020.5.3.8.7

According to microbial data, it seems that acetoclastic methanogens is the key pathway for methane production. The gas analysis of produced biogas showed CH_4 63 \pm 11, CO_2 35 \pm 9%, and

 H_2S 1800 \pm 105 ppm. Presence of *Syntrophomonas* and *Anaerobaculum* in the digester help to have more effective SAO.




Figure 5: Proposed pathways for CH₄ production in the studied digester; here, SAB and AM shown by solid grey arrows. SAO and HM pathways shown by dashed arrows. "Abbreviations: SAB: syntrophic acetogenic bacteria; SAOB: syntrophic acetate oxidizing bacteria; AM: acetoclastic methanogens; HM: hydrogenotrophic methanogens."

Conclusion

The sequencing study of this work was able to separate the microbial community and pathway of methane production in the full-scale AD of Tehran WWTP, successfully. According to our findings, 15 phyla were prevailing in the sludge of anaerobic digester, but Bacteroidetes and Firmicutes were the most abundance. the chief However, recognized genera were Anaerobaculum, Acinetobacter, Syntrophomonas, and Coprothermobacter, respectively. Exploring the leading bacteria and field data revealed that hydrogenotrophic methanogenesis acetate was metabolized by SAOBs. Generally, the findings of present study help the wastewater operators and process designers to use most effective method that able to treat waste sludge and methane production, simultaneously.

Acknowledgements

This study was supported by Tehran South wastewater treatment plant. We thank the WWTP management for helping us to conduct this study.

Funding

This study was financially supported by WWTP, south of Tehran, Iran.

Conflict of interest

The authors declare that they have no competing financial interests or personal relationships in this paper.

This is an Open-Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt, and build upon this work, for commercial use.

References

- 1.Hoseinzadeh E. Nitrogen removal processes for wastewater treatment: Bentham science publishers; 2019.
- 2.Scholz M. Chapter 15-Activated sludge processes. In: Scholz M, editor. Wetlands for water pollution control. Elsevier; 2016.
- 3.Song J, Chen L, Chen H, et al. Characterization and high-throughput sequencing of a trichlorophenol-dechlorinating microbial community acclimated from sewage sludge. J Clean Prod. 2018;197:306-13.
- 4.Canziani R, Spinosa L. 1-Sludge from wastewater treatment plants. In: Prasad MNV, de Campos Favas PJ, Vithanage M, et al., editors. Industrial

- and Municipal Sludge: Butterworth-Heinemann; 2019: 3-30.
- 5. Amani T, Nosrati M, Sreekrishnan T. Anaerobic digestion from the viewpoint of microbiological, chemical, and operational aspects-a review. Environmental Reviews. 2010;18:255-78.
- 6.Mir MA, Hussain A, Verma C. Design considerations and operational performance of anaerobic digester: A review. Cogent Eng. 2016;3(1):1181696.
- 7.Morales-Polo C, Cledera-Castro MDM, Moratilla Soria BY. Reviewing the anaerobic digestion of food waste: From waste generation and anaerobic process to its perspectives. Appl Sci. 2018;8(10):1804.
- 8. Kleerebezem R, Joosse B, Rozendal R, et al. Anaerobic digestion without biogas? Rev Environ Sci Biotechnol. 2015;14(4):787-801.
- 9.Guo J, Peng Y, Ni B-J, et al. Dissecting microbial community structure and methane-producing pathways of a full-scale anaerobic reactor digesting activated sludge from wastewater treatment by metagenomic sequencing. Microb Cell Fact. 2015;14:33.
- 10. Vanwonterghem I, Jensen PD, Ho DP, et al. Linking microbial community structure, interactions and function in anaerobic digesters using new molecular techniques. Curr Opin Biotechnol. 2014;27:55-64.
- 11. Luo C, Tsementzi D, Kyrpides N, et al. Direct comparisons of illumina vs. roche 454 sequencing technologies on the same microbial community DNA sample. PloS One. 2012;7(2):e30087.
- 12. Reuter JA, Spacek DV, Snyder MP. High-throughput sequencing technologies. Molecular Cell. 2015;58(4):586-97.
- 13. Singer E, Bushnell B, Coleman-Derr D, et al. High-resolution phylogenetic microbial community profiling. ISME J. 2016;10:2020.
- 14. Chao Y, Ma L, Yang Y, et al. Metagenomic analysis reveals significant changes of microbial compositions and protective functions during drinking water treatment. Scientific Reports. 2013;3:3550.
- 15. Chao Y, Mao Y, Wang Z, et al. Diversity and functions of bacterial community in drinking

- water biofilms revealed by high-throughput sequencing. Scientific Reports. 2015;5:10044.
- 16. Ding Y, Liang Z, Guo Z, et al. The performance and microbial community identification in mesophilic and atmospheric anaerobic membrane bioreactor for municipal wastewater treatment associated with different hydraulic retention times. Water. 2019;11(1): 160.
- 17. Kim B-R, Shin J, Guevarra R, et al. Deciphering diversity indices for a better understanding of microbial communities. J Microbiol Biotechnol. 2017;27(12):2089-93.
- 18. Kundu K, Sharma S, Sreekrishnan TR. Changes in microbial communities in a hybrid anaerobic reactor with organic loading rate and temperature. Bioresour Technol. 2013;129: 538-47.
- 19. Chao A, Chazdon RL, Colwell RK, et al. Abundance-based similarity indices and their estimation when there are unseen species in samples. Biometrics. 2006;62(2):361-71.
- 20. Musa MA, Idrus S, Hasfalina CM, et al. Effect of organic loading rate on anaerobic digestion performance of mesophilic (UASB) reactor using cattle slaughterhouse wastewater as substrate. Int J Environ Res Public Health. 2018;15(10):2220.
- 21. Aoyagi T, Hamai T, Hori T, et al. Hydraulic retention time and pH affect the performance and microbial communities of passive bioreactors for treatment of acid mine drainage. AMB Express. 2017;7(1):142.
- 22. ElNaker NA, Yousef AF, Hasan SW. Effect of hydraulic retention time on microbial community structure in wastewater treatment electro-bioreactors. Microbiologyopen. 2018;7(4): e00590.

Jehsd.ssu.ac.ir

- 23. Lim JW, Ge T, Tong YW. Monitoring of microbial communities in anaerobic digestion sludge for biogas optimisation. Waste Manag. 2018;71:334-41.
- 24. Li J, Rui J, Pei Z, et al. Straw- and slurry-associated prokaryotic communities differ during co-fermentation of straw and swine manure. Appl Microbiol Biotechnol. 2014; 98(10):4771-80.
- 25. Zhang B, Xu X, Zhu L. Structure and function of the microbial consortia of activated sludge in

- typical municipal wastewater treatment plants in winter. Scientific Reports. 2017;7(1):17930.
- 26. Rios-Covian D, Cuesta I, Alvarez-Buylla JR, et al. Bacteroides fragilis metabolises exopolysaccharides produced by bifidobacteria. BMC Microbiol. 2016;16(1):150.
- 27. Narihiro T, Kim N-K, Mei R, et al. Microbial community analysis of anaerobic reactors treating soft drink wastewater. PLoS One. 2015;10(3): e0119131.
- 28. Cai M, Wilkins D, Chen J, et al. Metagenomic reconstruction of key anaerobic digestion pathways in municipal sludge and industrial wastewater biogas-producing systems. Front Microbiol. 2016;7:778.
- 29. Nascimento AL, Souza AJ, Andrade PAM, et al. Sewage sludge microbial structures and relations to their sources, treatments, and chemical attributes. Front Microbiol. 2018;9:1462.
- 30. Martín-Díaz J, Ruiz-Hernando M, Astals S, et al. Assessing the usefulness of clostridia spores for evaluating sewage sludge hygienization. Bioresour Technol. 2017;225:286-92.
- 31. Pradhan N, Dipasquale L, d'Ippolito G, et al. Hydrogen production by the thermophilic bacterium thermotoga neapolitana. Int J Mol Sci. 2015;16(6):12578-600.
- 32. Lefèvre CT, Viloria N, Schmidt ML, et al. Novel magnetite-producing magnetotactic bacteria belonging to the gammaproteobacteria. ISME J. 2012;6(2):440-50.
- 33. Hoseinzadeh E, Rezaee A, Farzadkia M. Enhanced biological nitrate removal by alternating electric current bioelectrical reactor: Selectivity and mechanism. J Mol Liq. 2017;246:93-102.

- 34. Hoseinzadeh E, Rezaee A, Farzadkia M. Nitrate removal from pharmaceutical wastewater using microbial electrochemical system supplied through low frequency-low voltage alternating electric current. J Bioelectrochemistry. 2018;120: 49-56.
- 35. Hoseinzadeh E, Rezaee A, Fazeli S. Electrochemical denitrification using carbon cloth as an efficient anode. Desalination Water Treat. 2017;97:244-50.
- 36. Ushiki N, Fujitani H, Aoi Y, et al. Isolation of nitrospira belonging to sublineage II from a wastewater treatment plant. Microbes Environ. 2013;28(3):346-53.
- 37. Maune MW, Tanner RS. Description of Anaerobaculum hydrogeniformans sp. nov., an anaerobe that produces hydrogen from glucose, and emended description of the genus Anaerobaculum. Int J Syst Evol Microbiol. 2012; 62(4):832-8.
- 38. Sugihara T, Shiratori H, Domoto R, et al. Unique diversity content of a thermophilic anaerobic microbial consortium that utilizes propionate in a synthetic medium. J Gen Appl Microbiol. 2007;53(6): 363-9.
- 39. Esteves EMM, Herrera AMN, Esteves VPP, et al. Life cycle assessment of manure biogas production: A review. J Clean Prod. 2019;219:411-23.
- 40. Mosbæk F, Kjeldal H, Mulat DG, et al. Identification of syntrophic acetate-oxidizing bacteria in anaerobic digesters by combined protein-based stable isotope probing and metagenomics. ISME J. 2016;10(10): 2405-18.