

Journal of Environmental Health and Sustainable Development



Tardigrades in Wastewater Biomonitoring: Microscopic Organisms with Exceptional Environmental Resilience

Behnam Hatami 1, Mohammad Mehranpour 1*

¹ Environmental Sciences and Technology Research Center, Department of Environmental Health Engineering, School of Public Health, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.

ARTICLE INFO

LETTER TO EDITOR

Article History:

Received: 15 February 2025 Accepted: 20 April 2025

*Corresponding Author:

Mohammad Mehranpour

Fmail:

Mohammadmehranpor7@gmail.com

Tel:

+98 35 3149 2299

Citation: Hatami B, Mehranpour M. *Tardigrades in Wastewater Biomonitoring: Microscopic Organisms with Exceptional Environmental Resilience*. J Environ Health Sustain Dev. 2025; 10(2): 2586-8.

Tardigrades are microscopic, multicellular invertebrates with a cosmopolitan distribution across terrestrial and aquatic ecosystems. Due to physiological exceptional adaptations, particularly their capacity to endure extreme environmental stressors, they have emerged as model organisms in comparative physiology and extremophile biology. To date, over 1200 species have been documented from a broad range of ecological niches, including bryophytes, lichens, soil substrates, and both ephemeral and permanent aquatic habitats. A defining characteristic of tardigrades is their ability to undergo reversible ametabolic cryptobiosis—a state triggered by adverse environmental conditions such as severe desiccation, extreme temperature fluctuations ranging from 90 °C to -196 °C, hyperbaric pressures (up to 7.5 GPa), exposure to organic solvents, and ionizing radiation¹. During cryptobiosis, metabolic activity is virtually suspended, and the organism contracts into a desiccated, barrel-shaped form known as a "tun," wherein the cellular water content is reduced to as little as 2-3%, and body volume diminishes by approximately 85-90%. This anhydrobiotic state

facilitates survival in environments that exceed the tolerances of most known metazoans. Notably, tardigrades have demonstrated viability even after a 10-day exposure to the vacuum and radiation of outer space²⁻⁴.

In recent years, tardigrades have received increasing attention as potential bioindicators for assessing environmental quality in contaminated ecosystems, particularly within wastewater treatment facilities, owing to their exceptional physiological resilience under harsh environmental conditions. Species such as Thulinius ruffoi, observed in activated sludge, demonstrate the capacity to persist in environments characterized by high nutrient loads and complex chemical pollutants. The elevated abundance of such taxa during early stages of wastewater treatmenttypically associated with peak pollutant concentrations—underscores their potential utility monitoring effluent quality dynamics⁵. Nevertheless, the relatively limited distribution and low population densities of tardigrades in certain wastewater habitats, in comparison with more commonly employed indicator taxa such as protozoans and rotifers, have posed constraints on

Jehsd.ssu.ac.ir

[DOI: 10.18502/jehsd.v10i2.19004]

their broader application as routine bioassessment tools^{6, 7}. Consequently, while tardigrades may serve as complementary bioindicators within integrated environmental monitoring frameworks, their effective utilization necessitates the development of standardized sampling protocols and the execution of comprehensive studies to elucidate species-specific sensitivities to environmental perturbations^{5, 6}.

Given the extraordinary stress tolerance of tardigrades and emerging findings-such as the determination of LC50 values for Macrobiotus hufelandi upon lead exposure (94.651 mg/L at 24 hours and 8.048 mg/L at 96 hours)—demonstrating the necessity for expanded research into their environmental and biomedical applications has become increasingly evident8. The discovery of unique proteins such as the unique damage suppressor protein (Dsup), which binds to nucleosomes and protects DNA from hydroxyl radical-induced damage, and its demonstrated efficacy in enhancing human cell resistance to ionizing radiation and oxidative stress, has opened avenues for engineering stress-tolerant microbial strains⁹. The heterologous expression of tardigrade-derived genes in target microorganisms offers the potential to develop innovative bioremediation systems capable of maintaining pollutant-degrading functionality under extreme contamination conditions. Furthermore, leveraging tardigrades or their molecular biomarkers for monitoring water and soil quality—due to their pronounced tolerance to toxicants-may significantly enhance the sensitivity and reliability of environmental surveillance tools.

A particularly promising application involves the genetic simulation and transfer of tardigradespecific genes into beneficial bacteria used in wastewater treatment processes, with the aim of increasing their resistance to toxins and environmental stressors. This strategy could improve the performance of treatment systems operating in industrial or heavily polluted settings. addition. investigating the resistance mechanisms of tardigrades to various wastewater contaminants—such as heavy

pharmaceutical compounds—may yield insights into their survival strategies under extreme conditions and inform the development of next-generation wastewater treatment technologies. However, current challenges related to the large-scale production of these proteins underscore the need for advances in synthetic biology platforms and optimization of gene expression using state-of-the-art biotechnological methods. Achieving these objectives will require integrative approaches that synergize molecular biology, genetic engineering, and environmental science^{2, 8}.

This is an Open-Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt, and build upon this work for commercial use.

References

- 1. Zarubin M, Azorskaya T, Kuldoshina O, et al. The tardigrade Dsup protein enhances radioresistance in Drosophila melanogaster and acts as an unspecific repressor of transcription. Iscience. 2023;26(7).
- Kasianchuk N, Rzymski P, Kaczmarek Ł. The biomedical potential of tardigrade proteins: a review. Biomedicine & Pharmacotherapy. 2023; 158:114063.
- 3. Møbjerg N, Neves RC. New insights into survival strategies of tardigrades. Comp Biochem Physiol A Mol Integr Physiol. 2021;254:110890.
- 4. Jönsson KI, Rabbow E, Schill RO, et al. Tardigrades survive exposure to space in low Earth orbit. Current biology. 2008;18(17):R729-R31.
- 5. Jakubowska-Krepska N, Goldyn B, Krzemińska-Wowk P, et al. Tardigrades as potential bioindicators in biological wastewater treatment plants. Eur J Ecol. 2018;4(2): 124-30.
- Sobczyk M, Michno K, Kosztyła P, et al. Tolerance to ammonia of Thulinius ruffoi (Bertolani, 1981), a tardigrade isolated from a sewage treatment plant. Bull Environ Contam Toxicol. 2015;95:721-7.
- 7. Vargha B, Otvos E, Tuba Z. Investigations on ecological effects of heavy metal pollution in

2588

- Hungary by moss-dwelling water bears [Tardigrada], as bioindicators. Ann Agric Environ Med. 2002;9(2).
- 8. Berdi D, Nassouhi D, Çiftci E, et al. Acute toxicity of lead on the survival of Macrobiotus hufelandi (Eutardigrada: Parachela: Macrobiotidae). J Wildl
- Biodivers. 2024;8(2):38-54.
- 9. Chavez C, Cruz-Becerra G, Fei J, et al. The tardigrade damage suppressor protein binds to nucleosomes and protects DNA from hydroxyl radicals. Elife. 2019;8:e47682.